

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number
WO 03/001197 A1

(51) International Patent Classification⁷: G01N 33/00,
A01K 67/00, 67/033, C12N 5/00, 5/02, 5/06, 5/10

Nashville, TN 37209 (US). MILLER, David [US/US];
909 South Lane Court, Brentwood, TN 37027 (US).

(21) International Application Number: PCT/US02/19230

(74) Agent: SUBRAMONY, Priya, D.; FULBRIGHT &
JAWORSKI L.L.P., 600 Congress Avenue, Suite 2400,
Austin, TX 78701 (US).

(22) International Filing Date: 18 June 2002 (18.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/888,233 22 June 2001 (22.06.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 09/888,233 (CON)
Filed on 22 June 2001 (22.06.2001)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): VAN-
DERBILT UNIVERSITY [US/US]; Box 6009 Station B,
Nashville, TN 37235 (US).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BLAKELY, Randy,
D. [US/US]; 1727 Reines Court, Brentwood, TN 37202
(US). NASS, Richard [US/US]; 541 Burges Drive,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ASSAY FOR TOXIN INDUCED NEURONAL DEGENERATION AND VIABILITY INC. *ELEGANS*

(57) Abstract: Provided are in vivo screening methods to detect and identify substances that affect neuronal viability, and/or prevent neurodegeneration, and/or confer neuroprotective effects. The screening methods utilize recombinant *C. elegans* expressing a detectable marker in neuronal sub/groups and the use of neurotoxins specific to specific neuronal cells. Also provided are methods for identifying modulators of neurotransmitter transporters such as the dopamine transporter. Therefore, the invention provides methods for identifying substances that can be used in the prevention and therapy of neurodegenerative diseases.

WO 03/001197 A1

DESCRIPTION

ASSAY FOR TOXIN INDUCED NEURONAL DEGENERATION AND VIABILITY IN *C. ELEGANS*

5

BACKGROUND OF THE INVENTION

The government owns rights in the present invention pursuant to grant number DK58212 from the National Institutes of Health.

10

1. Field of the Invention

The present invention relates generally to the fields of neurobiology and neurodegenerative diseases. More particularly, it concerns the development of screening methods to detect and identify candidate molecules that confer neuroprotective effects following toxin-induced neuronal damage. The invention describes *in vivo* screening protocols using recombinant *C. elegans* that expresses a detectable marker in sub-groups of neurons, such as, dopaminergic neurons.

15

2. Description of Related Art

Neurodegenerative diseases, strokes and neuronal injuries caused by trauma are typically characterized by neuronal cell death of groups of neurons. Among these neurological disorders, neurodegenerative diseases such as Alzheimer's, Parkinson's, Huntington's, amyotrophic lateral sclerosis, *etc.*, afflict a large percentage of the population. For example, Alzheimer's disease alone afflicts about 4 million people in the United States, primarily the elderly, and is characterized by progressive memory loss, disorientation, depression and eventual loss of other body functions. Amyotrophic lateral sclerosis afflicts about 30,000 Americans, typically begins after age 40 and results in progressive weakness and paralysis. Huntington's Disease afflicts an estimated 25,000 patients in the United States, usually begins between the ages of 30 and 50 and includes violent, involuntary movements. Parkinson's Disease (PD) affects over 1 million people in the United States, and also usually begins on or after age 50. PD is another progressive disorder of the central nervous system and is characterized by a decrease in spontaneous movements, gait difficulty, postural instability, rigidity and tremor due to the degeneration of dopaminergic neurons.

20

25

30

In spite of great effort, little is known about the molecular basis of these disorders. Although the use of vertebrate and tissue culture systems continue to provide valuable insight into the pathology of the neurodegeneration, the molecular determinants involved in the etiology of these diseases remain elusive. As a result, there is an acute deficiency of effective therapeutic agents to treat these neurodegenerative disorders.

Although several drugs currently are being used for treatment of these diseases, none of these drugs offer complete cure or reversal of these disorders. In fact, most drugs only temporarily relieve some of the symptoms associated with the disease and do not prevent further degeneration of neurons. Hence, these disorders have been termed as progressive neurological disorders. Thus, the main goal of researchers in this field is to identify agents that will provide preventive as well as therapeutic relief for such diseases. At present, there is no effective high-throughput method to identify molecules with neuroprotective abilities *in vivo*.

SUMMARY OF THE INVENTION

The present invention overcomes deficiencies in the art and provides effective screening methods to identify agents that confer protective effects on neurons. The subject invention is based on a screening assay developed using the microscopic transparent roundworm *Caenorhabditis elegans* (*C. elegans*), which is a particularly useful model for studying neurodegeneration because it allows observation of changes in cells within the living worm over the time-period that it takes to develop from a single-cell zygote to a mature adult. These kinds of observations are extremely difficult in other animals and impossible in humans. As the genetics of *C. elegans* are well known, and as the entire genomic sequence has been mapped, the *C. elegans* based screening system of the invention provides a powerful screening/identifying tool.

Therefore, one embodiment of the invention provides a recombinant *C. elegans* that expresses a detectable marker in a dopamine neuronal. In one aspect, the detectable marker is further defined as a marker that can be visually detected. In a particular aspect the detectable marker is further defined as a marker that can be spectroscopically detected. Some examples of detectable markers include green fluorescent proteins, yellow fluorescent proteins, blue fluorescent proteins and red fluorescent proteins. Other spectral variants, mutational variants

and derivatives of such fluorescent proteins are also contemplated. Fluorescent proteins are preferred as they permit *in vivo* visualization of cells of the live worm.

In another aspect the detectable marker is β -galactosidase. Use of a marker such as β -galactosidase is however limited as it does not permit the visualization of live worms. Methods for detecting the expression of β -galactosidase using the substrate X-Gal are known to the skilled artisan. In yet another aspect, the detectable marker is an antigenic polypeptide positioned under the control of a promoter which can then be detected by staining with a labeled antibody. Methods for detection using antibodies are well known in the art.

In one embodiment the detectable marker is under the control of a promoter. In specific embodiments the promoter is a tissue-specific promoter. In other specific embodiments, the tissue-specific promoter is a neuronal promoter. Some neuronal promoters contemplated as useful are promoters specific to dopamine neurons such as the dopamine transporter promoters and the tyrosine hydroxylase promoters. In one embodiment, the dopamine transporter promoter region comprises the CeDAT promoter region comprised in SEQ. ID. NO. 1.

Expression of the detectable marker driven by specific promoters allows for the selective expression of the marker in specific subsets of cells, such as in dopamine neurons, cholinergic motor neurons, *etc.* The skilled artisan will recognize that the present invention is not limited to the use of the promoters described above and that any other specific neuronal, or other tissue specific promoter may be used in the practice of this invention. The main goal is to obtain expression of a detectable marker in certain specific populations of cells.

The strains of *C. elegans* provided by the subject invention can also be used in methods of screening drugs and other agents to identify substances that can prevent or decrease neuronal degeneration. In some aspects the neuronal degeneration is induced by a neurotoxic agent. One of the *C. elegans* strain is designated RN200 and comprises a *C. elegans* in which the dopamine transporter promoter drives the expression of a green fluorescent protein in a dopamine neuron.

Therefore, the invention further provides methods for screening for substances that affect neuronal viability comprising: (a) providing a recombinant *C. elegans* that expresses a detectable marker in a neuronal cell; (b) exposing the *C. elegans* to a candidate substance; and (c) detecting a change in the expression of the marker relative to the expression of the marker

before the exposing, wherein a change in the expression of the marker corresponds to a change in the viability of the neuron. The method may further comprise detecting the expression of the marker in the neuronal cell in the absence of the candidate substance. In one embodiment, the method further comprises the step of exposing the *C. elegans* to a known neurotoxin prior to step (b).

The candidate substance may be a substance that affects neuronal viability by either decreasing neuronal viability or by increasing neuronal viability. The screening is intended to identify both neuroprotective substances as well as neurotoxic substances.

Therefore, in some aspects, the substance is a neurotoxic substance. In specific aspects, the neurotoxic substance is 6-hydroxydopamine, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, or 5,7-di-hydroxy tryptamine. In other specific aspects, the neurotoxic substance is a generator of free radical species. In still other aspects, the substance is an environmental toxin. Examples of environmental toxins include pesticides, herbicides, air pollutants, water pollutants, industrial wastes. The term "pesticide" as used herein refers to any type of chemical compound which has lethal or sub-lethal effects on eukaryotic cells. Encompassed within "pesticide" are such compounds as insecticides, vermicides, mutagens, carcinogens, and any other compound useful for killing, mutating, or debilitating organisms such as insects. "Pesticides" include, but are not limited to polychlorinated hydrocarbons such as dichlorodiphenyltrichloroethane (DDT), dieldrin, aldrin, chlordane and lindane; organophosphorous compounds; carbamates; rotenone; and any other compound useful in the killing of pests.

In other aspects, the substance is a neuroprotective substance. In specific aspects, the neuroprotective substance is a dopamine transporter antagonist. In other specific aspects the dopamine transporter antagonist is a *C. elegans* specific dopamine transporter antagonist (Jayanthi *et al.*, 1998). Some non-limiting examples of dopamine transporter antagonists contemplated include imipramine, ampetamines, and cocaine.

In yet other aspects, the neuroprotective substance is a free radical scavenger. In specific embodiments, free radical scavengers include ascorbic acid (vitamin C), vitamin E, melatonin, and carboxyfullerenes, as some non-limiting examples. It also is contemplated that the substance may be encoded by a polynucleotide. In one embodiment, it is contemplated that the polynucleotide encodes a dopamine transporter regulatory gene, or a gene that suppresses free

radical generation. In other embodiments it is contemplated that the substance is a polypeptide. The polypeptide may encode a dopamine transporter regulatory polypeptide, or a polypeptide that suppresses free radical generation.

5 In other embodiments, the substance is a naturally occurring product. In yet other embodiments, the substance can be a man-made chemical. Some examples of substances contemplated as useful include monoamine oxidase (MAO) inhibitors. MAO inhibitors have been recently used in clinical trials for limiting the progress of PD. MAO inhibitors include hydrazine derivatives such as phenelzine or isocarboxazid or non-hydrazine derivatives such as
10 tranylcypromine, or pargyline.

Therefore it is contemplated that any synthetic compound, or natural product, including macromolecular entities such as polypeptides, polynucleotides, or lipids and also small entities such as neurotransmitters, ligands, hormones or elemental compounds may be used as candidate
15 substances. The synthetic compounds or natural products further refer substances that are either part of a crude mixture or purified and isolated, or just naturally occurring.

Other aspects of the method further comprise: (a) exposing the *C. elegans* to a known neurotoxin; and (b) detecting a change in expression of the marker. The change in marker
20 expression can be an increase in the marker or a decrease in the marker expression.

In one aspect, the detectable marker is further defined as a marker that can be visually detected. In another aspect, the detectable marker is further defined as a marker that can be spectroscopically detected. Some examples of detectable markers include green fluorescent
25 proteins, yellow fluorescent proteins, blue fluorescent proteins and red fluorescent proteins. Other spectral variants, mutational variants and derivatives of such fluorescent proteins are also contemplated. Use of fluorescent proteins is a preferred embodiment as it allows the screening in live animals which represents the most intact physiology and permits high-throughput *in vivo* screening.

30 In another aspect the detectable marker is β -galactosidase. Methods for detecting the expression of β -galactosidase using the substrate X-Gal are known to the skilled artisan. In yet another aspect the detectable marker is an antigenic polypeptide positioned under the control of a

promoter which can then be detected by staining with a labeled antibody. Methods for detection using antibodies are well known in the art.

5 In one embodiment the detectable marker is under the control of a promoter. In specific
embodiments the promoter is a tissue-specific promoter. In other specific embodiments the
tissue-specific promoter is a neuronal promoter. Some neuronal promoters contemplated as
useful, are dopamine transporter promoters, tyrosine hydroxylase promoters (encoded by the *cat-*
2 gene), the *cha-1* promoter which is predominantly expressed in cholinergic neurons, the *acr-2*
10 promoter which is predominantly expressed in cholinergic motor neurons, the *unc-30* promoter
which is expressed by predominantly in GABA-ergic neurons, the *unc-4* promoter which is
expressed by A-class motoneurons. Expression of the detectable marker under specific
promoters allows for the selective expression of the marker in specific subsets of cells, such as in
dopamine neurons, cholinergic motor neurons, *etc.* The skilled artisan will recognize that the
15 present invention is not limited to the use of the promoters described above and that any other
specific neuronal, or other tissue specific promoter may be used in the practice of this invention.
The main goal is to obtain expression of a detectable marker in certain specific populations of
cells.

20 In one embodiment of the method, the neuronal cell comprises a dopaminergic neuron, a
cholinergic neuron, a GABA-ergic neuron, a glycinergic neuron, a serotonergic neuron, a
cholinergic motor neuron, a glutamatergic neuron, or a peptidergic neuron.

25 Other non-neuronal cellular populations also are contemplated for cell specific
expression of a detectable marker and include, vulval cells, pharyngeal cells, and excretory cells,
among others.

30 Additionally, provided are methods of screening for substances that can inhibit neuronal
cell death comprising: (a) providing a recombinant *C. elegans* that expresses a detectable marker
in a neuronal cell; (b) exposing the *C. elegans* to a known neurotoxin and a candidate substance;
(c) detecting expression of the marker; and (d) comparing the expression of the marker to the
expression of the marker in the absence of the candidate substance. In some embodiments of this
method, the *C. elegans* is exposed to the neurotoxin prior to the candidate substance. In other
embodiments of this method, the *C. elegans* is exposed to the candidate substance prior to the
neurotoxin.

The invention also provides methods of screening candidate substances to identify a substance that can be used for prevention and/or therapy of neurodegenerative diseases comprising: (a) obtaining a recombinant *C. elegans* that expresses a detectable marker in a neuronal cell under the control of a neuronal-specific promoter; (b) exposing the *C. elegans* to a known neurotoxin and a candidate substance; (c) detecting expression of the marker; and (d) comparing the expression of the marker to the expression of the marker in the absence of the candidate substance. The *C. elegans* may be exposed to the neurotoxin prior to the candidate substance or alternatively prior to the neurotoxin.

Neurodegenerative diseases contemplated include non-limiting examples such as, Parkinson's disease, Alzheimer's disease, Huntington's disease, a transmissible spongiform encephalopathy, a familial amyloid polyneuropathy (FAP), a prion diseases, a Tauopathy, a Trinucleotide disease, amyotrophic lateral sclerosis (ALS) or multiple system atrophy.

The invention also provides methods of screening for substances that modulate dopamine transporter function comprising: (a) obtaining a recombinant *C. elegans* that expresses a detectable marker in a dopaminergic neuronal cell; (b) exposing the *C. elegans* to a candidate substance; (c) exposing the *C. elegans* to a neurotoxin that requires a dopamine transporter for intracellular access; and (d) detecting any change in the expression of the GFP after step c.

In one embodiment, the candidate substance modulates the expression of the dopamine transporter. Modulation can be at the level of transcription, translation, post-translational modifications, insertion into the neuronal membrane, *etc.*

In another embodiment of this method, the candidate substance blocks transport by the dopamine transporter. In other embodiments, the candidate substance increases transport by the dopamine transporter. In yet another embodiment, the candidate substance is a addictive substance. Non-limiting examples of such addictive substances include cocaine, amphetamines, or methylphenidate. In still another embodiment, the candidate substance is a modulator of regulatory pathways that control the dopamine transporter promoter.

In one aspect, the method can be used to identify substances that provide therapy for neurological diseases involving dopamine transporter function. Examples of neurological

diseases involving dopamine transporter function include schizophrenia, addiction disorders, attention deficit hyperactivity disorder (ADHD), psychoses, Tourette's syndrome, or Parkinson's disease.

5 The invention also provides a method of screening for molecules that modulate presynaptic neuronal signaling comprising: (a) obtaining a recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a knockout for a component of neuronal signaling; (b) obtaining a second recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a mutant for a component of neuronal signaling; (c)
10 comparing the differences in neuronal viability when exposed to a neurotoxic substance in the *C. elegans* of step a) with the *C. elegans* of step (b); and (d) identifying the genetic component of the mutation. In one embodiment the method further comprises isolating the genetic component of neuronal signaling. In specific aspects the component of neuronal signaling can be signaling molecules that control dopamine metabolism such as α -methyl-paratyrosine, reserpine, or
15 tetrabenazine.

 Further contemplated is a method for screening post-mortem human or other animal brains for substances that can cause neurotoxicity. For example, it is contemplated that one would screen for substances, from a post-mortem brain of a patient with a neurodegenerative
20 disease, that cause neurotoxicity in *C. elegans* neurons using the screening methods of the present invention. It is contemplated that initially tissue/cellular extracts, fluids, and similar partially purified preparations will be screened. The substances may be then isolated and identified. It is envisioned that one can then develop diagnostic assays to diagnose a subject for the presence of such neurotoxic substances thereby allowing one to predict the possibility of a
25 neurodegenerative disease.

 As used in this specification, "a" or "an" mean one or more. As used in the instant claim(s), when used in conjunction with the word "comprising", the words "a" or "an" means one or more than one and the word "another" means at least a second or more.

30 Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit

and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIGS. 1A & 1B. A 3D confocal reconstruction (FIG. 1A) and differential interference contrast (DIC) image (FIG. 1B) of the six dopaminergic DA neurons located in the head of a hermaphrodite expressing green fluorescent protein (GFP) driven by the endogenous *C. elegans* dopamine transporter (CeDAT) promotor. Arrows identify the CEP and ADE DA neurons.

FIG. 2. Genetic screens to identify CeDAT mutants, CeDAT regulators, and determinants of toxin sensitivity using ethylmethanesulfonate (EMS).

FIGS. 3A, 3B, 3C & 3D. Direct visualization of DA neurons in living *C. elegans*. (FIG. 3A), CEP and ADE neurons and processes as revealed by confocal 3D reconstruction of GFP signal in the anterior end of animals transgenic for a PDAT1::GFP transcriptional reporter. (FIG. 3B), Schematic illustration of the neurons shown in FIG. 1A. (FIG. 3C), PDE neurons and processes as revealed by confocal 3D reconstruction of GFP signal in the posterior-lateral segment of animals transgenic for a PDAT1::GFP transcriptional reporter. (FIG. 3D), Schematic illustration of the neurons shown in FIG. 3A. In FIG. 3A and FIG. 3C, large full arrow designates cell bodies whereas other arrows point to processes of the appropriate cell body. Scale bar= ~10 micrometers.

FIGS. 4A, 4B, 4C, 4D, 4E & 4F. Effect of 6-OHDA on GFP signal in DA Neurons. (FIG. 4A), Transgenic worms treated with vehicle (ascorbic acid) prior to visualization. (FIGS. 4B, 4C & 4D), Worms as in FIG. 4A but treated with 10 mM 6-OHDA for 1 hr prior to plating and visualized 72 hrs later via confocal microscopy-various stages of apparent degeneration are evident. (FIG. 4E), Transgenic worms with PDAT1::GFP reporter coinubated with 6-OHDA

and 10 mM amphetamine. (FIG. 4F), Transgenic worms with PDAT1::GFP reporter coincubated with 6-OHDA and 10 mM imipramine. Coinubcation with either amphetamine or imipramine prevents visible evidence of 6-OHDA toxicity, consistent with pharmacological blockade of DAT-1.

5

FIGS. 5A, 5B, 5C, 5D, & 5E. Dependence of apparent 6-OHDA toxicity on DAT-1. (FIG. 5A), Control reporter transgenic line exposed to vehicle. (FIG. 5B), Reporter transgenic line exposed to 50 mM 6-OHDA. (FIG. 5C), Reporter transgenic line produced in a dat-1 deleted strain, exposed to vehicle. Reporter expression is identical to that seen in transgenic with normal levels of DAT-1. (FIG. 5D), Reporter transgenic line produced in a dat-1 deleted strain, exposed to 6-OHDA. Notice a failure of toxin to effect loss of GFP signal. (FIG. 5E), Plot of neuronal sensitivity to 6-OHDA in wildtype (blue) or dat-1 mutant (red) lines as a function of time of exposure. Whereas the wildtype line shows a time dependent loss of CEP neurons, the dat-1 deletion strain is protected.

15

FIGS. 6A, 6B, 6C & 6D. Transfer of the DAT gene product sensitizes non-DA neurons to 6-OHDA. (FIG. 6A), ASI chemosensory neurons made transgenic for a PDAF7::GFP reporter to reveal these cells and their processes. (FIG. 6B), Worms in FIG. 6A treated with 6-OHDA. Notice little or no effect of the toxin on these cells that lack DAT-1. (FIG. 6C), A transgenic line where PDAF7::DAT1::GFP translational fusion is expressed in the ASI neurons, bringing DAT- protein to these cells. (FIG. 6D), Animals in C, exposed to 6-OHDA. Transfer of the DAT1 transporter to the ASI cells conferred 6-OHDA sensitivity illustrating that DAT-1 transfer may allow other cells to be ablated and screened for selective responses to neurotoxins

25

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Neurodegenerative diseases such as Parkinson's disease (PD), transmissible spongiform encephalopathies (TSEs), Alzheimer's disease (AD), familial amyloid polyneuropathy (FAP) and Huntington's disease (HD), Prion diseases, *etc.*, are characterized by degeneration and loss of neuronal cells. To date there has been no successful therapeutic agent that can provide reversal and/or prevention of degeneration.

30

Although vertebrate systems and tissue cultures are available for studying neurodegenerative conditions, they are relatively difficult to manipulate and the time course of such studies are generally very long. Because of the high conservation of genes and metabolic pathways between invertebrates and humans, as well as the availability of genetic strategies to identify novel proteins, protein interactions, and drug targets, genetic analysis using invertebrate model systems provides an enormous potential in deducing the factors involved in neuronal disease.

The present inventors have developed a novel screening method using the nematode *Caenorhabditis elegans* (*C. elegans*) for identifying substances that affect neuronal viability, neuronal cell-death, neuronal survival, and/or neuronal degeneration. *C. elegans* offers a system which is easy to manipulate genetically and in culture and numerous genes identified in *C. elegans* have mammalian and human counterparts with a high level of sequence homology.

The inventors have developed a recombinant *C. elegans* that selectively expresses a detectable marker in a set of neuronal cells, such as, in a dopaminergic neuronal cell population, in a cholinergic neuronal population, etc. Using such recombinant *C. elegans* that express detectable markers in one neuronal population, the inventors established conditions under which known cell-specific neurotoxins lead to a decrease or loss in the expression of the detectable marker. The inventors then developed a screening method whereby one can expose or contact the recombinant *C. elegans* with substances/candidate substances that can prevent/reduce damage by a neurotoxin, which is reflected by a decrease in loss of marker gene expression. For example, using 6-hydroxydopamine (6-OHDA), which is neurotoxic to dopaminergic (DA) neurons, the inventors demonstrated a decrease in a detectable marker gene, such as the green fluorescent protein (GFP), in worms selectively expressing such a detectable marker gene in dopaminergic neurons. The inventors then demonstrated that the decrease/loss of the detectable marker gene expression can be prevented or reduced by agents that block the dopamine transporter protein (DAT), which is responsible for transporting the 6-OHDA into DA neurons (see FIGS. 4A-F).

Thus, the present invention provides a powerful screening method whereby one can screen for agents that can prevent neuronal degeneration and/or neuronal loss. As neurodegenerative diseases are characterized by loss/degeneration of populations of neurons, this assay provides methods to identify substances that can be used as therapeutic agents. For

example, in PD there is a selective degeneration of DA neurons in the substantia nigra. Thus, screening for agents that can prevent/decrease DA neuronal degeneration will provide putative therapeutic agents for therapy and prevention of PD. The invention is however not limited to PD and one may screen for substances that prevent the loss/degeneration of any set/subset/population of neurons. It is contemplated that the method will allow the identification of substances that can prevent neuronal damage by different types of neurotoxic agents, including environmental toxins, free radicals, drugs, chemicals. In addition, the method is contemplated to provide an assay for substances that affect neuronal viability. Therefore, it allows the detection of yet unknown neurotoxic agents as well as yet unknown neuroprotective agents.

Additionally, the system also provides a method for screening for substances that modulate DAT transporter function. The *C. elegans* DAT (CeDAT) is highly conserved to the vertebrate and human DAT protein. The DAT protein is the target of addictive substances such as the psychoactive stimulants including cocaine, methylphenidate, *etc.* Thus, substances identified by such a screening method can help identify therapeutics for addiction disorders. Additionally, the method also provides the ability to identify therapeutic agents for neurological disorders involving the DAT. This includes schizophrenia, attention deficit hyperactivity disorder, psychoses, Tourette's syndrome, *etc.*

In addition, the inventors contemplate methods to screen for modulators of presynaptic neuronal signaling using recombinant *C. elegans* expressing detectable markers in selective populations of neurons comprising monitoring responses to cell-specific toxins. These methods comprise comparing knockout strains of *C. elegans* that are deficient for a component of neuronal signaling, such as the DAT gene, with strains that have mutations in neuronal signaling pathways. Comparison of such strains can reveal genetic components involved in modulation of neuronal signaling by identifying the nature of the mutation and eventually the gene involved. For example, the mutation may be in other DAT regulatory genes, and/or in toxin suppressive genes, *etc.* Other methods contemplated include genetic methods whereby one can cross a knockout DAT *C. elegans* strain with a phenotypic DAT mutant *C. elegans* strain. If the cross rescues the knockout, this indicates that the mutation in the mutant strain is in a synaptic modulator or a toxin suppressor type gene, *i.e.*, in a gene other than the DAT gene. One can then identify the mutation and isolate and identify the gene corresponding to the synaptic modulator or toxin suppressor. Although the discussion above describes DAT protein and its modulators as

an example, this method is contemplated as useful for screening any molecule that can modulate neuronal signaling. Thus, one may screen for modulators of acetylcholine-mediated signaling, serotonin-mediated signaling, glutamate-mediated signaling, GABA-mediated signaling and the like.

5

A. Parkinson's Disease as a Model for Neurodegenerative Diseases

This section describes Parkinson's disease (PD) in detail, however, the general description applies to any neurodegenerative disorder. PD is a slowly progressive, neurodegenerative disorder characterized by the irreversible loss of over 80% of the nigrostriatal dopaminergic neurons. Although the pathogenesis of the disease appears to be multifactorial, correlative evidence supports the role of oxidative stress and mitochondrial dysfunction (Yahr and Bergmann, 1986). PD patients have reduced levels of mitochondrial complex I and glutathione activity, as well as increased levels of superoxide dismutase activity, lipid peroxidation, and iron in the substantia nigra (Yahr and Bergmann, 1986; Jenner, 1998). This evidence suggests that the generation of reactive oxygen species (ROS), such as the superoxide ion and hydrogen peroxide plays a significant role in dopamine (DA) neuronal death.

Current animal model systems of PD rely on inducing nigrostriatal damage in monkeys and rodents with the neurotoxins 6-hydroxydopamine (6-OHDA), 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), or 1-methyl-4-phenylpyridinium ion (MPP⁺ [the active metabolite of MPTP]) (Yahr and Bergmann, 1986). Toxin exposure causes DA neuronal death, and the animals exhibit many of the same symptoms as PD. The specificity of the toxin for the DA neurons lies in their affinity for the Na⁺- and Cl⁻-dependent dopamine transporter (DAT), a presynaptic membrane protein that terminates DAergic transmission by the active reuptake of DA into presynaptic neurons (Sachs and Jonsson, 1975; Glinka *et al.*, 1997; Kitayama *et al.*, 1992; Pifl *et al.*, 1993; Javitch *et al.*, 1985; Giros *et al.*, 1996; Gainetdinov *et al.*, 1998). Transfections of DATs into nonneuronal cells confer sensitivity to MPP⁺, and 6-OHDA and MPP⁺ toxicity can be blocked with DAT antagonist *in vivo* (Glinka *et al.*, 1997; Kitayama *et al.*, 1992; Pifl *et al.*, 1993; Javitch *et al.*, 1985).

30

Despite decades of research, the primary insults and mechanisms leading to the degeneration of the nigrostriatal DA neurons and the increase in production of ROS in PD is unknown. Non-human primates do not naturally develop PD, so these model systems are unavailable to study the etiology of the disease. Current animal and tissue culture systems are

able to mimic some of the pathology and morphology of idiopathic PD. However, identifying the molecular determinants involved in PD without any *a priori* knowledge of the mechanism of the neurodegeneration is significantly hindered due to the lack of a definitive model system.

5 An approach to identifying the molecular determinants involved in PD without any prior knowledge of their function could be through forward genetics (Johnsen *et al.*, 1997). In this classical method of genetic analysis, a well-defined phenotype is modified by mutagenizing the genomic DNA. The progeny that have the modified trait are further analyzed to determine the location of the altered loci (this is the opposite of reverse genetics in which the goal is to identify
10 animals that have a specific gene mutated) (Johnsen *et al.*, 1997). In order for forward genetics to be successful in studying DAergic neurodegeneration in an animal, either an aspect of the behavior should depend on normal DA neurotransmission, or the morphology of the DA neurons should be easily accessible. Furthermore, the organism should provide an efficient strategy to map and identify the mutated gene. Because of the above criteria, as well as their relatively long
15 life-span and maintenance cost, mammals are not amenable for these studies.

C. elegans offers a powerful tool for dissecting the components involved in mammalian neurodegenerative disorders using forward genetics. *C. elegans* provides a model system due to numerous factors such as, the high conservation of the genome and metabolic pathways between
20 nematodes and vertebrates; the ease of screening a large number of worms following mutagenesis for neural defects; and the ability to identify quickly the modified gene as the whole genome of *C. elegans* been sequenced. Thus, the *C. elegans* model allows for the rapid discovery of proteins involved in neuropathologies (*C. elegans* Sequencing Consortium, 1998).

25 The rapid rate of reproduction of *C. elegans* (approximately 3 days at 25°C), and their relatively small size (approximately 1 mm), generates about 300 worms from a single hermaphrodite, allowing for rapid genetic screens involving large numbers of animals (Wood, 1988; Riddle *et al.*, 1997). Yet other features of *C. elegans* that make them amenable for studies are the transparency of the worm, with virtually every cell identifiable under a light microscope.
30 Furthermore, the entire 302-cell nervous system (approximately one-third of the total somatic cells of the hermaphrodite) has been mapped by serial 3-dimensional electron-microscopic reconstruction, allowing for the identification of virtually all synaptic connections (White *et al.*, 1986).

Multicolor reporters, using spectral derivatives of the green fluorescent protein (GFP) from the jellyfish *Aequorea victoria*, also allow for the detailed visualization of cells, intracellular compartments, or protein localization in the living animal (Chalfie *et al.*, 1994; Rongo *et al.*, 1998; Miller *et al.*, 1999b). Methods for generating transgenic animals (obtainable within 4 days) and production of gene knockouts averaging at about 1 week are also well developed (see Chapter 19, Mello and Fire, 1995). A significant advantage of the worm over vertebrate systems is that sexual reproduction can be achieved by self-fertilization and therefore is not disrupted by mutations that would otherwise perturb mating behavior.

The similarity between the worm and the mammalian nervous system indicates that the fundamental interactions that occur during DA neurotransmission and pathogenesis in vertebrates also occurs in *C. elegans*. Most of the molecular components known to be involved in DA signaling in mammals are also present in the nematode (Table 1) (Loer and Kenyon, 1993; Lints and Emmons, 1999; Duerr *et al.*, 1999; Jayanthi *et al.*, 1998; Sulston *et al.*, 1992). Furthermore, analysis of *C. elegans* genome has revealed a high conservation of ion channels, neurotransmitter synthesis enzymes, including tyrosine hydroxylase (TH) and aromatic amino acid decarboxylase (AAAD); synaptic vesicle and presynaptic terminal proteins, including syntaxins, synaptotagmin, synaptobrevin; neurotransmitter receptors, including glutamate, acetylcholine, GABA, amine, and peptide; and neurotransmitter transporters, including acetylcholine (VchAT), the vesicular monoamine transporter (VMAT), GABA, glutamate (EAAT), serotonin (SERT), dopamine (DAT) (Table 1) (Bargmann, 1998).

Table 1
Genes of Catecholamine Biosynthesis, Metabolism, and Transport in *C. elegans*

Gene	Protein	Reference
<i>cat-4</i>	GTP Cyclohydrolase I	Loer and Kenyon, 1993
<i>cat-2</i>	Tyrosine Hydroxylase	Lints and Emmons, 1999
<i>bas-1</i>	Aromatic L-Amino Acid Decarboxylase	Loer and Kenyon, 1993
<i>cat-1</i>	Vesicular Monoamine Transporter	Duerr <i>et al.</i> , 1999
<i>dat-1</i>	Dopamine Transporter	Jayanthi <i>et al.</i> , 1998
R13G10.2	Monoamine Oxidase*	Sulston <i>et al.</i> , 1992
C48D5.1	NHR, Nurr1 Homologue*	BLAST search at http://www.wormbase.org/perl/ace/elegans/searches/blast

*Predicted gene product; NHR, nuclear hormone receptor

In addition, a mammalian nuclear hormone receptor, Nurr1, which is expressed in DA neurons and provides a measure of neuroprotection against MPTP toxicity, also has significant

homology with the worm gene C48D5.1 (Baffi *et al.*, 1999; Le *et al.*, 1999; Castillo *et al.*, 1998; www.wormbase.org/perl/ace/elegans/searches/blast). However to date, the cellular expression pattern of C48D5.1 has not been defined. An important distinction to make between the two nervous systems at the molecular level is the redundancy of the genomes, while vertebrates will often have several similar versions of a particular gene, *C. elegans* may have a single allele (Bargmann, 1998). This feature of the nematode genome provides an enormous advantage relative to vertebrate systems because it decreases the probability that a mutant effect will be masked by a genome redundancy. Finally, the genetic basis of cell death is highly conserved between the worm and vertebrates, and *C. elegans* provides an opportunity to explore genes involved in neuronal cell death (Hengartner, 1997).

The *C. elegans* hermaphrodite contains eight dopaminergic neurons which include six head DA neurons, four cephalic neurons (CEPs), and two anterior dereid neurons (ADEs), and two neurons which are located in posterior lateral positions, the posterior dereid neurons (PDEs) (Sulston *et al.*, 1975). All of these neurons contain ciliated endings which terminate in sensory organs embedded in the cuticle that surrounds the animal. Laser ablation and genetic studies have revealed that these cells provide mechanosensory function during foraging and movement and may also modulate pharyngeal pumping and egg laying behaviors (Duerr *et al.*, 1999; Sulston *et al.*, 1975; Sawin *et al.*, 2000; Lints and Emmons, 1999). The male contains another three pairs of DA-containing neurons in the tail that are involved in mating, as well as four additional DA-containing socket cells of the mating spicules (Loer and Kenyon, 1993; Lints and Emmons, 1999; Sulston *et al.*, 1975; Sulston and Horvitz, 1977; Liu and Sternberg, 1995).

B. Detectable Markers

The present invention contemplates the use of detectable markers that are selectively expressed in a specific set or population of neurons. Thus, the marker can be expressed, for example, selectively in DA neurons and not in others.

In the case of *C. elegans*, as the organism is transparent, one may also be able to detect the colored products produced by such marker genes directly by the human eye. One major class of detectable marker genes that may be used to visualize a live *C. elegans* worm in its intact form comprise the fluorescent proteins such as the green fluorescent proteins (GFP). GFP is a naturally fluorescent protein which can be used to mark the cells in which a promoter is active. It has the benefit that the animals can be observed live.

The GFP proteins, originally isolated from the jellyfish *Aequorea victoria* retain their fluorescent properties when expressed in heterologous cells thereby provides a powerful tool as fluorescent recombinant probes to monitor cellular events or functions (Chalfie *et al.* 1994; Prasher 1995; WO 95/07463). Chalfie *et al.*, in U.S. Patent 6,146,826, (incorporated herein by reference), also describes the expression of GFP in *C. elegans* touch receptor neurons under the control of a promoter for the *mec-7* gene which encodes a β -tubulin.

Several spectral and mutational variants of GFP proteins have since been isolated, for example, the naturally occurring blue-fluorescent variant of GFP (Heim *et al.* 1994; U.S. Patent 6,172,188, both incorporated herein by reference), the yellow-fluorescent protein variant of GFP (Miller *et al.*, 1999), and more recently the red fluorescent protein isolated from the coral *Discosoma* (Frédérkov *et al.*, 2000; Miller *et al.*, 1999), which allows the use of fluorescent probes having different excitation and emission spectra permitting the simultaneous monitoring of more than one process. GFP proteins provide non-invasive assays which allow detection of cellular events in intact, living cells. The skilled artisan will recognize that the invention is not limited to the fluorescent proteins described and one may use any other spectral variant or derivative.

It is contemplated that one or more detectable markers may be used to monitor the viability of one or more neuronal populations simultaneously. For example, if the organism is subject to an free radical generator, or a environmental toxin, one can label DA neurons with one marker, the cholinergic neurons with another marker, the serotonergic neurons with yet another marker and study the effects of such a toxin on all the three neuronal populations simultaneously. In some embodiments, it is contemplated that the use of multiple markers may contribute to high-throughput screening.

Other detectable markers are also contemplated as useful. These include, β -galactosidase, enzymes such as urease, alkaline phosphatase, horseradish peroxidase, which can be detected either spectrophotometrically by the use of colorimetric indicator substrates that are typically employed with these detectable genes. These markers are somewhat limited in their use as they cannot be used to monitor live worms and worms must be killed and fixed to visualize detection.

In the case of β -galactosidase, lacZ is a stable enzyme, with negligible background activity in *C. elegans*. The chromogenic stain is stable, and the enzyme is still active after mild fixation. Generally, a lacZ with a nuclear localization signal, derived from SV40, is used so that the nuclei of cells expressing the construct can be identified.

5

It is also possible to attach antigenic polypeptides or short tags of amino acids to the native genes, and then detect expression patterns by using an antibody which is specific to the antigenic polypeptide or tag.

10 Detectable markers are generally known in the art and the skilled artisan will recognize that the invention is not limited to the examples described above.

C. Transporters and Neurotransmission

Chemical signaling by small molecule neurotransmitters, including glutamate, glycine, GABA, DA, NE, and 5HT, is terminated by transporter-mediated clearance (Rudnick and Clark, 1993). Pharmacologic or genetic disruption of transporter function elevates extracellular neurotransmitter levels, perturbs presynaptic transmitter homeostasis, and can trigger significant alterations in behavior (Giros *et al.*, 1996; Pelham, 1997). The psychoactive agents cocaine and the amphetamines compete with substrates at amine (DA, NE, 5HT) transporters, with much of their addictive potential attributed to DAT blockade (Kuhar *et al.*, 1991). In contrast, NE and 5HT transporter (NET and SERT, respectively) antagonists such as imipramine, desipramine, fluoxetine, and sertraline are important agents in the treatment of mood disorders, particularly depression (Barker and Blakely, 1995). The cloning of a rat GAT1 (Guastella *et al.*, 1990) and a human NET (Pacholczyk *et al.*, 1991) established the presence of a Na and Cl dependent transporter gene family whose members include transporters for most small neurotransmitters. The present inventors have previously cloned and characterized the *C. elegans* dopamine transporter gene, the *T23G5.5* gene which encodes the CeDAT protein, and found that CeDAT is sensitive to antidepressants and cocaine (Jayanthi *et al.*, 1998).

30 (i) Dopaminergic Transmission

Dopamine (DA), is a biogenic amine synthesized in the hypothalamus, in the arcuate nucleus, the caudate, and in other areas of the central and peripheral nervous system. Dopamine is also a precursor of other neurotransmitters, specifically, norepinephrine (NE) and epinephrine (E), in addition to being a neurotransmitter on its own. Dopamine and its agonists play

important roles in cardiovascular, renal, hormonal, and central, nervous system regulation through stimulation of alpha and beta adrenergic and dopaminergic receptors.

DA being a catechol and easily oxidized to a quinone, is often implicated as a generator of reactive oxygen species (ROS) like peroxide (H_2O_2), superoxide (O_2^-) and hydroxyl radical (OH^\cdot) the latter being the most reactive and detrimental ROS. The dopamine transporter protein (DAT) is responsible for the uptake of excess dopamine that is released into the synaptic space back into neurons. Uptake of dopamine by DAT is important for regulating neuronal signaling as well as reducing the potential for DA being oxidized to form ROS.

(ii) Dopamine Transporter Protein (DAT)

DAT is a plasma membrane transport protein that controls extracellular DA concentrations, by recapture of DA that has been released during the process of neurotransmission, into nerve terminal. More recently, DAT has been recognized as a major target for various pharmacologically active drugs and environmental toxins (Miller, *et al.*, 1999b).

DAT has been cloned and information regarding its structure and function are available (see review, by Chen, 2000, and the references cited therein). Chen (2000), describes that binding domains for dopamine and various blocking drugs including cocaine are likely formed by interactions with multiple amino acid residues, some of which are separate in the primary structure but lie close together in the still unknown tertiary structure. Other studies utilizing chimeric proteins and site-directed mutagenesis suggest the involvement of both overlapping and separate domains in the interaction with substrates and blockers. Recent findings using sulfhydryl reagents that selectively target cysteine residues support a role for conformational changes in the binding of DAT antagonists such as cocaine. The dopamine transporter can also operate in reverse, *i.e.*, in an efflux mode, and recent mutagenesis experiments show different structural requirements for inward and outward transport.

DAT is also involved in various neurological and neuropsychiatric disorders ranging from Parkinson's disease to attention deficit disorder, schizophrenia, Tourette's syndrome, to drug abuse and addiction.

a. Sequence And Structure

DAT is a member of a large family of Na^+/Cl^- dependent transporters, which includes the norepinephrine (NE) transporter (NET) as well as transporters for serotonin, GABA, glycine, proline, creatine, betaine, taurine. Using sequences conserved between the GABA transporter and NET, DAT cDNAs have been cloned from rat, cow and human sources. DAT is a 619 (rat) or 620 (human) amino acid protein and the hydropathy and immunochemical data analysis indicate that DAT includes 12 transmembrane domains (TMDs), with both the amino- and carboxy-termini residing within the cytoplasm.

The generation of chimeric transporters produced by exchanging similar domains between the highly related DAT and NET transporters has provided clues as to the functional aspects of different DAT domains. It appears that residues within TMDs 1-3, or alternatively TMDs 9-12, greatly influence substrate affinities. TMDs 1-3 also influence affinity for certain DAT ligands such as GBR12395. TMDs 5-8 may also be critical in determining affinities for DAT and NET inhibitors such as cocaine and desipramine. The affinity of the DA neurotoxin MPP+ for the DAT is influenced by residues within TMDs 11-12.

b. DAT Gene Expression

The DAT gene is expressed largely in the CNS of mammals within a small subset of DA neurons and not in glia. As DAT expression is more restricted, than the expression of other genes such as the genes encoding DA biosynthetic enzymes such as, tyrosine hydroxylase (TH), aromatic amino acid decarboxylase, or the DA receptors, DAT provides an excellent marker for DA neurons and their projections.

In the rodent, DAT mRNA is found in great abundance within midbrain DA neurons of the substantia nigra, with somewhat lower expression in the ventral tegmental nuclei and adjacent nuclei. Within the hypothalamus, DAT is expressed within the A13 (zona incerta) and, to a lesser extent, the A14 (periventricular) and A12 (arcuate nucleus) cell groups, but not other TH-positive cell groups. Moderate DAT expression is also seen in the A16 cell group of the olfactory bulb. DAT mRNA is not found in regions devoid of DA cell bodies or within DA nerve terminals. In human brain, DAT mRNA exhibits the same general distribution as seen in rodents. Within human midbrain DA cells, the abundance of DAT mRNA is greatest within the ventral tier of the substantia nigra, followed by the dorsal tier and the ventral tegmental area, with the lowest levels of DAT mRNA seen within the retrorubral field.

The distribution of DAT immunoreactivity is largely consistent with other indices of DAT gene expression and the density of DA innervation. Thus, the striatum and nucleus accumbens are densely labeled, with labeling also apparent within the globus pallidus, cingulate cortex, olfactory tubercle and amygdala. DAT immunoreactivity is also seen in the perikarya, dendrites and axonal processes of midbrain DA neurons. Regional differences in somatodendritic DAT immunoreactivity covary with DAT mRNA levels and correlate with the susceptibility of subgroups of DA neurons to neurotoxins and idiopathic disease processes. DAT protein is localized primarily on extrasynaptic plasma membranes near aggregates of synaptic vesicles, consistent with DAT playing a key role in limiting the spatial domain of DA neurotransmission. Within the dendrites of midbrain DA cells, DAT is localized to plasma membranes and smooth endoplasmic reticulum, consistent with DAT-mediated dendritic DA release and/or DAT modulation of DA cell activity through its channel-like properties. Within perikarya, DAT is localized primarily to tubulovesicles, which may represent DA-containing membranes in transit.

c. DAT Gene Organization

The hDAT gene has been localized to chromosome 5p15.3 (35; 108), cloned and characterized. The gene spans 64 kb and is divided into 15 exons separated by 14 introns, with predicted intron-exon junctions. The hDAT coding region begins within exon 2 and extends partially into exon 15. A single transcriptional start site has been identified. There is no evidence for DAT RNA splice variants or the use of multiple polyadenylation sites. The overall exon-intron structure of the hDAT gene closely parallels that of the hNET (and to a lesser extent the serotonin and GABA transporter) gene. In general, each hDAT exon encodes a functional domain such as the N- or C-terminus or a putative TMD and adjacent hydrophilic loop. A number of Alu and other sequence repeats have been located within intronic portions of the gene. Of greater interest is a 40 bp variable tandem nucleotide repeat (VNTR) polymorphic sequence found in the 3' untranslated region just upstream of the polyadenylation site which may be associated with human diseases. A number of restriction fragment length polymorphisms have also been examined for such associations.

The 5' flanking sequences controlling transcription of the hDAT gene are interesting in that neither a canonical 'TATA' box nor a 'CAAT' box have been found, prompting the suggestion that the DAT is a TATA-less gene. The high local GC content and several putative

SP1 sites might serve to direct DAT gene transcription. Other than the potential SP1 sites, a rather limited number of potential transcription factor response elements (such as, Egr-1, E-box, AP-2) may also comprise the proximal hDAT promoter, although individual elements have not been assessed functionally.

5

Recently, over 8 kb of hDAT 5' flanking sequence has been cloned (GenBank accession Number AF115382, incorporated herein by reference). Within this span of sequence, numerous potential regulatory sequences have been tentatively identified but, as with the core promoter, no genomic element that affords cell specific DA neuron expression has been identified. The development of midbrain DA cells is critically dependent on the expression of the nuclear receptor transcription factor *nurr1*, as proven by targeted disruption ("knockout") of the *nurr1* gene. The *nurr1* gene expression persists in rodent midbrain DA cells through adulthood, suggesting post-developmental functions as well. Interestingly, multiple *nurr1* binding sites have been identified in 5' flanking sequences of the hDAT gene, and hDAT promoter constructs are activated by *nurr1* co-transfection *in vitro*. Furthermore, the human homologue of *nurr1* (NOT-1) is expressed at high levels in human midbrain DA neurons. Hypothalamic DA neurons express much lower levels of both *nurr1* and DAT. Thus, it appears that the nuclear receptor *nurr1*/NOT-1 plays an important role in the maintenance, as well as development, of the strong DAergic phenotype seen in midbrain.

20

d. Psychostimulants and DAT Knockout Animals

DAT is a major target for psychostimulants such as cocaine, amphetamine and methamphetamine. The reinforcing properties of these drugs have been strongly correlated with their affinities for the DAT, in keeping with evidence implicating mesolimbic DA neurons in drug abuse. Cocaine and related drugs bind to the DAT and prevent DA transport. Amphetamines gain access to DA nerve terminals by both lipophilic diffusion across the plasma membrane and DAT-mediated transport, releasing DA from vesicular stores and evoking DAT-mediated DA release. In each instance, psychostimulants raise the extracellular concentrations of DA, likely augmenting DA neurotransmission over longer distances and/or durations, and causing behavioral activation.

30

Confirmation of these facts has been obtained by examination of DAT gene knockout animals. These mice exhibit significant locomotor hyperactivity in response to cocaine or amphetamine. In the absence of DAT, DA clearance time from the extracellular space is

markedly prolonged (100-300 fold) and stimulants fail to further augment extracellular DA concentrations. It is interesting that this phenotype is accompanied by only 5 fold increases in extracellular DA, presumably due to profound (75-95%) decreases in the DA biosynthetic enzyme TH, DA levels, and DA release. Reduced expression of basal ganglia D1 and D2 DA receptors is also evident. In spite of low basal expression of DAT in the hypothalamus, the development of anterior pituitary hypoplasia and dwarfism in DAT knockouts suggests an unanticipated importance of the DAT in regulating pituitary function. The knockout mice are resistant to the DA neurotoxic effects of MPP+ and methamphetamine.

e. Potential Role of DAT in Human Disorders

DA neurotransmission has been implicated in various neuropsychiatric disorders, including Parkinson's disease, schizophrenia, Tourette's syndrome, attention-deficit disorder, and substance abuse. DAT is the primary DA-binding protein that functions to remove the neurotransmitter from the synaptic space. Given the nonredundancy of the gene, and its central role in controlling spatial and temporal aspects of DA neurotransmission, the DAT gene has received considerable attention as a candidate gene for DA-related neuropsychiatric disorders.

As mentioned above, characterization of the hDAT cDNA and gene has identified a VNTR. Although, the role of VNTRs in gene function remains obscure, recent evidence suggests that these repetitive elements could play a role in transcriptional and post-transcriptional gene expression. There are currently no data indicating that the VNTR in the DAT gene affects its expression. Nevertheless, the polymorphic VNTR serves as a highly informative marker for association and linkage analyses.

The DAT VNTR is 40 bp sequence that is present in the 3' nontranslated region of the DAT cDNA and is repeated 3-11 times. The most common allele contains 10 copies of the VNTR. In the US population, the frequency of the allele is 0.7 among Caucasians and Hispanics, and about 0.54 in African Americans. Among Asians, the frequency of the 10 copy allele is about 0.9, making the degree of heterozygosity in these populations quite low. The degree of ethnic heterogeneity in the DAT VNTR is an important consideration in interpreting disease association studies.

Perhaps the best evidence for involvement of the DAT with a disorder comes from work with attention-deficit hyperactivity disorder (ADHD). ADHD appears to be familial and

heritable, and is perhaps the most common childhood-onset behavioral disorder. It is well recognized that ADHD patients benefit from treatment with certain psychostimulants, such as methylphenidate and amphetamine, which directly interact with the DAT. The evaluation of DAT as a candidate for susceptibility to ADHD using haplotype-based haplotype relative risk analysis. This study found a significant association of the 10 copy VNTR polymorphism with ADHD and undifferentiated ADHD. Similar results were independently obtained in an Irish population.

The DAT also has been proposed as a gene candidate for Parkinson's disease owing to the potential ability of DAT to transport neurotoxins into the DA neurons that are destroyed in the disease (Miller, 1999b).

DAT in Parkinson's Disease. PD is characterized by a substantial loss of midbrain DA neurons with a consequent loss of DA innervation to forebrain structures. The vulnerability of certain subgroups of DA neurons in PD and MPTP-induced parkinsonism correlates with higher basal levels of DAT gene expression. It is conceivable that avid transport of neurotoxins or even endogenous DA by the DAT may play a role in idiopathic PD.

Given the extent of DA cell loss, it is not surprising that significant decreases in DAT ligand binding sites are detected in PD striatum postmortem. DAT binding is reduced equivalently in progressive supranuclear palsy, a disease involving global degenerative changes throughout the basal ganglia and associated nuclei including the substantia nigra.

DAT in Alzheimer's Disease with Parkinsonism. Signs of clinical Parkinsonism occur in a sizable proportion (20-40%) of patients with Alzheimer's disease in the absence of classical neuropathology of PD. Alzheimer's subjects with parkinsonism exhibit a substantial loss of DAT in the caudate-putamen, albeit with a distribution which differs from the loss of DAT in idiopathic PD. There is also a significant decline in the number of DAT mRNA-positive DA cells in the midbrain, as well as lower DAT mRNA levels per cell. The profound loss of midbrain DA cells which occurs in PD, however, is not seen in parkinsonian Alzheimer's subjects, and TH expression is less impacted than the expression of DAT. DA-related gene expression, therefore, is impacted differently in Alzheimer's disease with parkinsonism than in idiopathic PD, although the underlying mechanisms are not understood. In nonparkinsonian Alzheimer's subjects, DAT and DA systems in general are minimally impacted.

DAT in Wilson's Disease. Wilson's disease is an autosomal recessive disorder involving mutations of the P-type copper ATPase ATP7B, resulting in excess copper deposition. Liver and brain are most affected. Varied neurological symptoms include parkinsonism, dystonia, and psychosis. Structural changes are seen in numerous brain regions including the striatum, where a loss of D2 receptors occurs. DAT ligand binding *in vivo* is decreased in Wilson's disease to the same extent as seen in PD, but without corresponding pathological changes in the substantia nigra. These preliminary data are somewhat reminiscent of the changes seen in Alzheimer's disease with parkinsonism and suggest a preferential loss of DA nerve terminals or a profound change in DAT biosynthesis, transport, or turnover.

DAT in Lesch-Nyhan Disease. This X-linked infantile onset disease, resulting from the loss of hypoxanthine-guanine phosphoribosyl transferase activity, leads to compulsive self-injurious behavior and movement disorders such as dystonia and choreoathetosis. Self-injurious behavior can be elicited in rodents by neonatal manipulations of DA function, and DA systems seem to be significantly impacted in Lesch-Nyhan. Striatal DAT density determined by PET imaging reported to be decreased 50-75% in a small group of Lesch-Nyhan subjects; other measures of DA function and total striatal volume are also significantly affected in this severe disorder.

DAT in Tourette's Syndrome. This disorder is characterized by symptoms including obsessions, compulsions, coprolalia and involuntary tics. Although the neuropathological mechanisms underlying Tourette's syndrome are unknown, the therapeutic benefit of decreasing DA neurotransmission is cited as evidence for the involvement of DA systems. A provocative study reported 37-50% increases in the density of DAT ligand binding in postmortem caudate-putamen from a small number of Tourette's syndrome subjects compared to controls. The concentrations of striatal DA and DA metabolites and DA receptor density were reportedly unchanged. More recently, a SPECT imaging study involving a small number of Tourette's subjects reported an equivalent (37%) increase in DAT density. It is speculated that an undetermined perturbation of DA systems in Tourette's syndrome may lead to a compensatory upregulation of striatal DAT.

DAT in Schizophrenia. The longstanding suspicion that DA systems are somehow involved in the etiology, symptomatology and/or treatment response of schizophrenia has led to

investigations of DAT expression in this disorder. A functional study reported alterations in the V_{max} and K_m for DA uptake into cryopreserved nerve terminals from the striatum of schizophrenics relative to age- and sex-matched controls.

5 *DAT in Chronic Stimulant Abusers.* As discussed above, the DAT represents a major target protein for cocaine and amphetamine-like drugs. Chronic exposure to these drugs might therefore be expected to elicit compensatory changes in DAT expression. Thus, DAT is strongly implicated in drug addiction disorders.

10 (iii) CeDAT

Jayanthi *et al.* (1998) report the cloning and characterization of the *C. elegans* DAT (CeDAT). CeDAT is highly conserved with the mammalian catecholamine transporters with an amino acid identity of approximately 45%. The gene encodes a 615 amino-acid protein with 12 putative transmembrane domains. Mammalian cells expressing CeDAT exhibit saturable and
15 high affinity Na^+ - and Cl^- -dependent DA transport ($K_m=1.2 \mu M$) and are sensitive to the mammalian DAT antagonist GBR 12909 and cocaine, and to the substrate d-amphetamine at potencies comparable to their actions on human DAT in transfected cells (Jayanthi *et al.*, 1998; Giros *et al.*, 1992; Giros and Caron, 1993). CeDAT is also inhibited by the tricyclic antidepressant SERT antagonist imipramine and the NET-selective antagonist nisoxetine
20 (Jayanthi *et al.*, 1998). Mammalian DAT proteins are exclusively expressed in dopamine neurons.

Kyte-Doolittle hydrophobicity analysis suggested twelve hydrophobic stretches suitable for formation of TMDs that are well aligned with similar profiles of transporters in the *gat1/net*
25 gene family. Two canonical sites for N-linked glycosylation are located in the large hydrophilic loop between TMDs 3 and 4, sites analogous to those known to be glycosylated in mammalian catecholamine transporters (Melikian *et al.*, 1996). Additional N-glycosylation sites are present in the transporter's amino (N22) and carboxyl (N597) termini. The amino and carboxyl termini possess a number of Ser and Thr residues that may be targets for regulatory phosphorylation with
30 two PKC sites (Ser45, Ser582) and one casein kinase II site (Thr580). A cAMP-dependent protein kinase site (Ser255) also lies in a putative intracellular loop between tmdsTMDs 4 and 5 within a span of residues (WKGXXTSKGKVVW) found in all catecholamine transporters. Similarly, a casein kinase II site between TMDs 6 and 7 lies in a highly conserved stretch of sequence (A(Y/F)SSYN(D/K)F. Comparisons with other *gat1/net* family members demonstrates

highest similarity of CeDAT to mammalian catecholamine transporters. CeDAT exhibits 47% amino acid identity with human, mouse, and bovine NETs, 43% identity with human, bovine, and rat DATs, 37% identity with human, rat and, mouse SERTs, and less than 35% identity with other gene family members. Sequence divergence suggests the carrier may have arisen from a common ancestral transporter before DATs, NETs, and SERTs formed genetically distinct species. An Asp residue that is conserved in TMD1 of the DA, NE and 5HT transporters from fly to man (Kitayama *et al.*, 1992; Barker and Blakely, 1995) but absent from GABA, glycine, taurine, proline, creatine and taurine transporters, is also present in an analogous position (D60) in CeDAT. Thus, there is sequence divergence and conservation that is evident comparing CeDAT with its most closely related mammalian homologs.

D. Expression Vectors

Within certain embodiments, expression vectors are employed to express a detectable marker, such as a GFP polypeptide, beta-galactosidase, or an antigenic polypeptide. Expression requires that appropriate signals be provided in the vectors, and which include various regulatory elements, such as enhancers/promoters from both viral and mammalian sources that drive expression of the genes of interest in host cells. Elements designed to optimize messenger RNA stability and translatability in host cells also are defined. The conditions for the use of a number of dominant drug selection markers for establishing permanent, stable cell clones expressing the products are also provided, as is an element that links expression of the drug selection markers to expression of the polypeptide.

Throughout this application, the term "expression construct" is meant to include any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed and translated into a polypeptide product. An "expression cassette" is defined as a nucleic acid encoding a gene product under transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrase "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

The term promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about

how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (*tk*) and SV40 early transcription units. These studies, augmented by more recent work, have shown that promoters are composed of discrete functional modules, each consisting of approximately 7-20 bp of DNA, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the *tk* promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

By attaching a tissue-specific or cell-specific promoter region of a nucleic acid to a reporter or a detectable marker, one can obtain tissue-specific or cell-specific expression of the nucleic acid of interest. *C. elegans* promoter elements tend to be relatively short, often less than 1 kb. Both transcriptional and translational fusions can be made.

To derive neuronal expression of marker genes some promoters contemplated useful in context of the present invention include the dopamine transporter promoter, the tyrosine hydroxylase promoter (encoded by the *cat-2* gene), the *cha-1* promoter (expressed in cholinergic neurons), an *acr-2* promoter (expressed in cholinergic motor neurons), an *unc-30* promoter (expressed by GABA-ergic neurons), or an *unc-4* promoter (expressed by A-class motoneurons).

In the case of the dopamine transporter promoter the present inventors have shown that a ~700bp region of the CeDAT promoter region is responsible for DA neuron specific reporter expression. The CeDAT promoter regions are also described in SEQ ID. NO. 1.

5 The use of other *C. elegans* promoters which are well-known in the art to achieve expression of a coding sequence of interest are contemplated as well, provided that the levels of expression are sufficient for a given purpose.

10 In addition, the use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression of a coding sequence of interest are contemplated as well.

15 By employing a promoter with well-known properties, the level and pattern of expression of the protein of interest following transfection or transformation can be optimized. Further, selection of a promoter that is regulated in response to specific physiologic signals can permit inducible expression of the gene product.

20 Enhancers are genetic elements that increase transcription from a promoter located at a distant position on the same molecule of DNA. Enhancers are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

25 The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

30 **E. Screening For Modulators Of Neuronal Viability and/or Signaling**

 The present invention comprises methods for screening for a variety of substances, including substances that affect neuronal viability; substances that inhibit neuronal cell death; substances that prevent neuronal degeneration; substances that cause neuronal degeneration, and/or substances that modulate neuronal signaling and function. The present invention further

comprises genetic methods for identifying modulators of neuronal signaling such as modulators of DAT, modulators of dopamine signaling and/or other elements that modulate presynaptic signaling.

5 These assays may comprise random screening of large libraries of candidate substances; alternatively, the assays may be used to focus on particular classes of compounds selected with an eye towards structural attributes that are believed to make them more likely to modulate the function of neuronal cells, such as known neurotoxins, neurotransmitter transporter agonists and antagonists, addictive substances, pesticides, therapeutic substances, monoamine oxidase
10 inhibitors, and antioxidants.

 By function, it is meant that one may assay for the expression of a detectable marker expressed selectively in some neuronal populations. A decrease or loss in the expression of the marker indicates neuronal cell death, apoptosis, degeneration, and/or decrease in neuronal
15 viability. On the other hand, no difference in the expression of the marker indicates that the cells were either protected from degeneration (if a second substance is used) or were not affected by the substance. The preferred detectable marker is a fluorescent protein as it allows for longitudinal studies on live animals, although other markers such as beta-galactosidase, enzymes that react to produce colored reaction products, may also be used.

20

 To screening for substances that, affect neuronal viability; and/or ability to prevent neuronal degeneration; and/or cause neuronal degeneration; and/or inhibit neuronal cell-death, a method will generally comprise: (a) providing a *C. elegans* that expresses a detectable marker in a neuronal cell; (b) exposing the *C. elegans* to a candidate substance; and (c) detecting a
25 change in the expression of the marker relative to the expression of the marker before said exposing; wherein a change in the expression of the marker corresponds to a change in the viability; and/or ability to prevent neuronal degeneration; and/or ability to cause neuronal degeneration; and/or ability to inhibit neuronal cell-death, of the neuron.

30

 To identify a modulator of dopamine transporter protein (DAT), one generally will determine the function of DAT in the presence and absence of the candidate substance. A modulator is defined as any substance that alters function. For example, a method generally comprises: (a) obtaining a recombinant *C. elegans* that expresses a detectable marker in a dopaminergic neuronal cell; (b) exposing said *C. elegans* to a candidate substance; (c) exposing

said *C. elegans* to a neurotoxin that requires a dopamine transporter for intracellular access; and (d) detecting any change in the expression of the GFP after step (c), wherein a difference between the measured characteristics, *i.e.*, expression of marker, indicates that said candidate modulator substance is, indeed, a modulator of the DAT.

5

The invention also describes genetic screening that is aimed at identifying the genetic component of the modulators of neuronal signaling. Such a method of screening for molecules that modulate neuronal signaling comprises: (a) obtaining a recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a knockout and/or a mutant for a component of neuronal signaling; (b) obtaining a second recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a (or another) mutant for a component of neuronal signaling; (c) comparing the differences in neuronal viability when exposed to a neurotoxic substance in the *C. elegans* of step (a) with the *C. elegans* of step (b); and (d) identifying the genetic component of the mutation.

10
15

Assays may be conducted in cell free systems, in isolated cells, or in organisms including transgenic animals, recombinant animals, mutated animals. A preferred embodiment uses live animals as this represents the most intact physiology and permits high-throughput screening.

20

It will, of course, be understood that all the screening methods of the present invention are useful in themselves notwithstanding the fact that effective candidates may not be found. The invention provides methods for screening for such candidates, not solely methods of finding them.

25

(i) Modulators

30

As used herein the term "candidate substance" refers to any molecule that may potentially inhibit or enhance DAT activity. The candidate substance may be a protein or fragment thereof, a small molecule, or even a nucleic acid molecule. It may prove to be the case that the most useful pharmacological compounds will be compounds that are structurally related to cocaine, methylphenidate, *etc.* Using lead compounds to help develop improved compounds is known as "rational drug design" and includes not only comparisons with known inhibitors and activators, but predictions relating to the structure of target molecules.

The goal of rational drug design is to produce structural analogs of biologically active polypeptides or target compounds. By creating such analogs, it is possible to fashion drugs, which are more active or stable than the natural molecules, which have different susceptibility to alteration or which may affect the function of various other molecules. In one approach, one
5 would generate a three-dimensional structure for a target molecule, or a fragment thereof. This could be accomplished by x-ray crystallography, computer modeling or by a combination of both approaches.

It also is possible to use antibodies to ascertain the structure of a target compound
10 activator or inhibitor. In principle, this approach yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of anti-idiotypic would be expected to be an analog of the original antigen. The anti-idiotypic could then be used to identify and isolate
15 peptides from banks of chemically- or biologically-produced peptides. Selected peptides would then serve as the pharmacore. Anti-idiotypes may be generated using the methods described herein for producing antibodies, using an antibody as the antigen.

On the other hand, one may simply acquire, from various commercial sources, small
20 molecule libraries that are believed to meet the basic criteria for useful drugs in an effort to "brute force" the identification of useful compounds. Screening of such libraries, including combinatorially generated libraries (e.g., peptide libraries), is a rapid and efficient way to screen large number of related (and unrelated) compounds for activity. Combinatorial approaches also lend themselves to rapid evolution of potential drugs by the creation of second, third and fourth
25 generation compounds modeled of active, but otherwise undesirable compounds.

Candidate compounds may include fragments or parts of naturally-occurring compounds, or may be found as active combinations of known compounds, which are otherwise inactive. It is proposed that compounds isolated from natural sources, such as animals, bacteria, fungi, plant
30 sources, including leaves and bark, and marine samples may be assayed as candidates for the presence of potentially useful pharmaceutical agents. It will be understood that the pharmaceutical agents to be screened could also be derived or synthesized from chemical compositions or man-made compounds. Thus, it is understood that the candidate substance identified by the present invention may be peptide, polypeptide, polynucleotide, small molecule

inhibitors or any other compounds that may be designed through rational drug design starting from known inhibitors or stimulators.

Other suitable modulators include antisense molecules, ribozymes, and antibodies (including single chain antibodies), each of which would be specific for the target molecule. Such compounds are described in greater detail elsewhere in this document. For example, an antisense molecule that bound to a translational or transcriptional start site, or splice junctions, would be ideal candidate inhibitors.

In addition to the modulating compounds initially identified, the inventors also contemplate that other sterically similar compounds may be formulated to mimic the key portions of the structure of the modulators. Such compounds, which may include peptidomimetics of peptide modulators, may be used in the same manner as the initial modulators.

An inhibitor according to the present invention may be one which exerts its inhibitory or activating effect upstream, downstream or directly on neuronal viability and neuronal signaling. Regardless of the type of inhibitor or activator identified by the present screening methods, the effect of the inhibition or activation by such a compound results in differences in neuronal viability or neuronal signaling as compared to that observed in the absence of the added candidate substance.

(ii) *In vivo* Assays

In vivo assays involve the use of various animal models, including transgenic animals that have been engineered to have specific defects, or carry markers that can be used to measure the ability of a candidate substance to reach and effect different cells within the organism. Due to their size, transparency, ease of handling, and information on their physiology and genetic make-up, *C. elegans* are a preferred embodiment.

In such assays, one or more candidate substances are administered to an animal, and the ability of the candidate substance(s) to alter one or more characteristics, as compared to a similar animal not treated with the candidate substance(s), identifies a modulator. The characteristics may be any of those discussed above with regard to the neuronal viability, inhibition of neuronal degeneration/cell death, or neuronal signaling.

Treatment of these animals with test compounds will involve the administration/exposing of the compound, in an appropriate form, to the animal. Administration or exposing of the compound to a worm is by adding the compound to a liquid suspension and transferring the worms to this suspension prior to replating the worms on solid media. More details are described in the section entitled as 'Examples'.

Determining the effectiveness of a compound *in vivo* may involve a variety of different criteria. Also, measuring toxicity and dose response can be performed in animals in a more meaningful fashion than in *in vitro* or *in cyto* assays.

(iii) Genetic Assays

Other genetic assays that comprising crossing a *C. elegans* that bears a mutation in a neuronal signaling component with a *C. elegans* that is a knockout for a neuronal signaling component may be performed. In such an assay, if the progeny of the cross demonstrates a rescue of the phenotype, this indicates that the mutation in the mutant strain is in a gene other than the gene of the knockout. One can then identify the mutation and isolate and identify the gene corresponding to the mutation using the powerful genetics of *C. elegans*.

a. Setting up Crosses

Although *C. elegans* is a self-fertilizing hermaphrodite it is possible to set up genetic crosses as functional males are found (albeit rarely) and male sperm outcompetes hermaphrodite sperm. Wild type *C. elegans* males are found at about 0.05% of the population. They develop after accidental non-disjunction of the X chromosomes in gametogenesis (males are XO). The rate of nondisjunction can be increased by exposing a population of hermaphrodites to heat shock (30°C for >6 hrs), and "male lines" can be propagated indefinitely by picking individual males and hermaphrodites. In addition, several loci have been described which when mutated result in high levels of males in a population. These Him strains, such as those carrying the him-8 mutation, can be used to generate males carrying desired mutations for crossing by mating with mutant hermaphrodites. When a hermaphrodite is mated, the male sperm outcompetes her own supply, and the progeny are thus almost entirely cross-fertilized.

F. Transgenic, Knockout and Mutant *C. elegans*

In an exemplary embodiment, the "transgenic animals" of the invention are produced by introducing transgenes into the germline of the of the animal. Different methods can be used for the introduction of transgenes. The specific line(s) of any animal used to practice this invention
5 are selected for general good health, good embryo yields, and good reproductive fitness.

The early germline of *C. elegans*, which are hermaphrodites, is syncytial, *i.e.*, individual nuclei reside in pockets of cytoplasm but are connected to a central rachis. This facilitates uptake of any nucleic acid injected into the syncytial gonad by multiple nuclei. However, the
10 germline nuclei that take up the nucleic acid, such as a DNA vector bearing the nucleic acid encoding a polypeptide of interest, have a repair machinery that catenates the nucleic acid into long extrachromosomal arrays (>100 kb) which are maintained in the resultant offspring with relatively stability. The transgenic nematodes are selected by co-injecting a visible marker gene, usually a dominant mutation in the cuticular collagen gene *rol-6(su1006)*. This gene causes the
15 nematodes to develop a helically twisted cuticle and to roll longitudinally when they move forward.

Transgenesis can be used for several sorts of analysis including as a proof that a DNA fragment contains the wild-type copy of a mutated gene, by rescuing the mutant with the
20 transgene. Transgenic animals maybe used to generate "antisense knockout" strains. In such embodiments, if the open reading frame of the nucleic acid is attached "backwards" to its own promoter, antisense RNA will be made, which suppress translation of the wild type mRNA. In other embodiments, analysis of promoter elements can be performed by ligating the 5' region of a nucleic acid to detectable marker genes, for example to the enzymatic markers such as *lacZ* or
25 beta-galactosidase, or to fluorescent reporters/markers, such as the green fluorescent proteins. Transgenic *C. elegans* may also be used to drive the overexpression of a nucleic acid to look at gain-of-function mutations in the gene encoded by the nucleic acid. One may also introduce *in vitro* mutagenized or foreign genes to examine structure-function relationships and to constructing complexly mutated strains.

30 Transgenic extrachromosomal arrays can be stably integrated into the chromosomes by UV irradiation of transgenic lines, or by coinjection of single stranded DNA. Homologous recombination is very rare in *C. elegans*.

The progeny of the transgenically manipulated worms can be tested for the presence of the construct by any of the following: 1) by observation of altered behavior due to the transgene; 2) by Southern blot analysis of a segment of tissue; 3) by PCR amplification of genomic DNA; and/or by co-injecting a fluorescent marker with the transgene and observing the retention of fluorescence of the transgenic marker in the DNA. If one or more copies of the exogenous cloned construct remains stably integrated into the genome of such transgenic *C. elegans*, it is possible to establish permanent transgenic lines carrying the transgenically added construct.

The litters of transgenically altered worms can be assayed after birth for the incorporation of the construct into the genome of the offspring. Preferably, this assay is accomplished by hybridizing a probe corresponding to the DNA sequence coding for the desired recombinant protein product or a segment thereof onto chromosomal material from the progeny. Those progeny found to contain at least one copy of the construct in their genome are grown to maturity.

The "knockout animals" of the invention can be produced either by generating transgenic animals as described above or by mutagenesis of the animal. Different mutagenesis methods can be used. The specific line(s) of any animal used to practice this invention are selected for general good health, good embryo yields, and good reproductive fitness. In addition, the haplotype is a significant factor.

The gene knockout strategy for generating knockout *C. Elegans* involves using a random mutagen, such as trimethylpsoralen, or EMS, to mutagenize a very large number of worms (see http://info.med.yale.edu/mbb/koelle/protocols/protocol_Gene_knockouts.html, *C. elegans* Gene Knockout Protocols, by Michael Koelle, Heather Hess, David Shechner). Briefly, the worms are divided into many small subcultures and allowed to have progeny. A portion of each subculture is stored alive in the freezer, and genomic DNA is made from the rest of the culture. Therefore, the DNA made from the siblings of the frozen worms and carries the same mutations as the frozen worms do.

At a very low frequency (~1/400,000 mutagenized genomes) the mutagenesis produces a small deletion (~200-4,000 bp) in any gene of interest to generate knockouts. At a high frequency the mutagenesis produces point mutants. For the knockouts, PCR primers flanking the gene of interest are used to amplify from the genomic DNA samples, deletions can be detected

because primer sites flanking such a deletion will be brought closer together and will generate a PCR amplicon smaller in size than that amplified from wild-type genomic DNA. When the PCR reaction is carried out under appropriate conditions small deletion amplicons are amplified much more efficiently than the larger wild-type amplicon. Thus DNA representing several thousand mutagenized genomes can be amplified in a single reaction and a deletion amplicon generated from just one of those genomes will still be detected on an ethidium bromide stained agarose gel. Once a DNA pool containing a deletion is thus identified, one can work back to identify the subculture of worms in which the deletion occurred, the frozen worms from that subculture can be thawed, and live animals carrying the deletion mutation can be identified.

G. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLE 1

Construction of Recombinant *C. elegans* Expressing Detectable Markers

Methods

Strains and Maintenance

C. elegans strains were cultured on either OP-50 or NA-22 at 22.5°C according to standard methods (Brenner, 1994). The N2 Bristol strain is the wild-type strain and can be obtained from the Caenorhabditis Genetics Center (University of Minnesota, Minneapolis, MN).

Plasmid Construction

The DAT-1 transcriptional GFP fusion (Pdat-1::GFP) was created by PCR amplification of the DAT-1 regulatory region in cosmid T23G5.5. The amplified product immediately upstream of the initiating codon ATG was digested with HindIII and BamHI, and the resultant 0.7 kb fragment was cloned into the HindIII and BamHI sites of the GFP reporter vector pPD95.73 (gift from Andy Fire), to create the construct called pRN200.

Germline Transformation

Transgenic animals were generated as previously described (Mello *et al.*, 1991; Mello and Fire, 1995). Transgenic animals containing the transcriptional fusion (pRN200) were
5 obtained after co-injection of 20 ng/ μ l of pRN200, 30 ng/ μ l of plasmid carrier DNA (pBluescript), and 50 ng/ μ l of pRF4 (rol-6(su1006)) (Kramer *et al.*, 1990) into the gonads of the N2 strain. Animals were cultured at 24.5°C, and F1 and F2 transgenic animals were selected by their roller phenotype. The transgenic array containing the dat-1 transcriptional fusion was then integrated into the genome using irradiation-based recombination. Two independent integrants
10 were isolated and both display similar GFP expression patterns; one integrant, RN200, was outcrossed four times and used for subsequent analysis (now called RN200x4). RN200x4 was mapped to within 2.4 MU from dpy-11 on chromosome V.

6-OHDA Assays

Synchronized cultures of nematodes were prepared by collecting gravid worms from one
15 or two 100 mm 8P/NA22 plates, washing them 2 times in approximately 10 mls dH₂O (2,000 rpm for 2 min), and incubating them in alkaline hypochlorite for 7-10 min. essentially as described (Emmons *et al.*, 1979; Epstein and Shakes, 1995). Following the 17-24 hour incubation in M9 to obtain the synchronized L1's, the worms were washed once in 10 mls
20 dH₂O, spread on 8P/NA22 plates and incubated at 24° C for 25-29 hours. The L3 larvae were collected, washed once or twice in dH₂O, and added to the assay mix (to a final OD600 worms of 0.1-0.2) containing 10 mM ascorbic acid containing 10 or 50 mM 6-hydroxydopamine +/- 10 mM d-amphetamine or 1 mM imipramine. The assay mix (1-2 mls) was incubated for 1 hour at 24° C, and mixed gently every 15 min. The worms were then washed in 0.5-1 ml dH₂O as above
25 and spread on NGM/OP50 plates. The worms were examined and scored 40-80 hours post-6-OHDA exposure on 2% agarose pads, using 2% NaN₃ to immobilize them. For electron microscopy, the worms are fixed approximately 60 hours following exposure to 6-OHDA.

Scoring of the CEPs were performed by applying approximately 50-60 worms on the
30 agar pad, and examined under a Zeiss M2 fluorescent dissecting scope. All 4 CEP dendrites were examined in each worm and the GFP fluorescence within the dendrites followed from the nerve ring to the tip of the nose. If any part of the dendrite was absent, the worm was considered to have altered CEPs. The DA neurons were examined in living animals using a Zeiss Axioskop with a planApo 40 and 63x objective and a Zeiss LSM410 confocal laser-scanning microscope.

Scoring of the CEPs were performed on a Zeiss M2 fluorescent dissecting scope using a 2X and 10X objective. For electron microscopy, worms were fixed in buffered (100 mM Hepes, pH 7.5) 3% glutaraldehyde, followed by postfixation in buffered 1 % OsO₄ (Hall, 1995; Sulston and Hodgkin, 1988). After encasement in 1% agar, samples were dehydrated and embedded in Polybed 812 resin (Polysciences). Serial sections were poststained in uranyl acetate followed by lead citrate.

EXAMPLE 2

Screening for Modulators of DAT

For the genetic and pharmacological evaluation of DAT and genes involved in altered DA neurotransmission, the *C. elegans* dopamine transporter (CeDAT) was cloned as described in Jayanthi *et al.*, 1998, incorporated herein by reference.

Following the initial observation that a genomic fragment comprising the CeDAT gene (2.1 kilobase pair fragment upstream of the start ATG to 0.7 kb downstream of the ATG) was sufficient to drive reporter expression in DA cell bodies (T. Ishihara, personal communication; See I. Hope, worm expression pattern for gene T23G5.5 (<http://bgypc086.leeds.ac.uk>), an integrated transgenic line expressing a CeDAT promotor-GFP fusion (P_{CeDAT}::GFP) containing 717 bp immediately upstream of the start ATG was created. This line displays intense GFP expression in all eight DA neurons within the hermaphrodite, but is not evident in other cells (FIG. 1). The axons and dendrites in these transgenic animals are clearly visible and can be easily observed as the animal moves under a fluorescent dissecting scope.

Screening. In order to set up a screen for molecules involved in DA neurodegeneration in *C. elegans*, a well-defined behavior dependent on the normal DAergic function is required. Alternatively, visualization of changes in morphology of the DA neurons can be used as a screening tool. As DA mediated behaviors are relatively subtle in *C. elegans* screening a large number of worms based on the modification of these behaviors could be tedious, a more efficient and robust screening method is to select for mutants within the transgenic lines expressing GFP that have altered DA neuronal morphology. Thus, GFP expressing *C. elegans* that are mutant in DA neuronal morphology were used.

Toxin Lesioning. The neurotoxins 6-OHDA and MPP⁺ provide cell-specific lesions reminiscent of the selective pathology of PD and are used as tools to probe the mechanism of environmentally triggered DAergic neural degeneration in animal models of PD (Yahr and Bergmann, 1986, Lotharius *et al.*, 1999; Miller *et al.*, 1999).

5

C. elegans DA neurons were also selectively degraded by 6-OHDA which enters the cells via CeDAT. Briefly, the worms are exposed to the neurotoxin (or any other candidate substance) dissolved in a liquid suspension prior to being replated on a solid plate.

10

Exposure to toxins is carried out by transferring the worms to a solution of water containing 10mM Ascorbic Acid and 10-50 mM 6OHDA in a final volume of 600μl. The worms are incubated for 1 hour at 22-24°C with mixing in silica coated tubes. The exposing is stopped by dilution with 1ml of water. The tube is then pulsed at 2000 rpm for 2 minutes, resuspended in water and the worms are spread on N6M plates (100mM plates) coated with OP50 *E.coli*, which the worms feed on.

15

Accordingly, a specific loss of the GFP expression in those neurons that expressed CeDAT was observed in the transgenic lines following exposure to the toxins. Based on this, a genetic screen for the retention of GFP in DA neurons following toxin exposure is contemplated.

20

Genetic Screening. The present inventors contemplate performing genetic screens based on the strategy described below. To establish the screen, the integrated transgenic line carrying the transcriptional reporter P_{CeDAT}::GFP will be used. Following mutagenesis of the last larval stage of the worm (*i.e.*, the L4 stage, where the maximum number of gametes are present in the hermaphrodite), the F2 generation (second generation of the mutagenized worm) will be subject to 6-OHDA or MPP⁺ and tested for loss of toxin sensitivity. As most mutations are recessive, screening in the F2 generation allows the mutations to become homozygous and therefore increases the likelihood of a modified phenotype. All mutants recovered will then be physically mapped using a two-factor linkage analysis, and the genetic map interval defined by three-factor crosses (Sulston and Hodgkin, 1988). Once the area of the mutated allele has been identified, candidate genes can be tested through microinjection of cosmid spanning the gene. The inventors further contemplate assaying whether DA neuron-specific expression of the wild-type gene rescues sensitivity to 6-OHDA. Finally, the impact of coexpression of CeDAT or

25

30

mammalian DAT with the identified protein or its mammalian homolog (if one has been identified) following toxin exposure in mammalian cells will be performed.

The genetic screen is also described in FIG. 2. Following mutagenesis and toxin exposure, any retention of GFP within a DA neuron is due to either the inability of the toxin to enter the cell (for example via CeDAT), suppression of some mediator of toxin sensitivity, or inhibition of neuronal degradative pathway(s).

Analysis of Mutants. The inventors contemplate that the methods described will identify CeDAT mutants that are regulators and modulators of CeDAT. Since DATs are required for toxin accumulation in mammalian DA neurons, the inventors expect to recover an allelic series of mutations in CeDAT which would modify its activity, as well as regulators and modulators of CeDAT. A variety of CeDAT mutants that limit the toxin's ability to enter the cell (and hence alter GFP expression), including those affecting transporter activity, stability, and localization, are also expected to be identified in the screen. These mutants would be especially valuable because the structural basis for DA transport and DAT regulation are poorly defined. Furthermore, the availability of DATs at the nerve terminals has been proposed to play a role in the vulnerability of DAergic neurons to PD, and this screen should assist in delineating its role in the neurodegeneration (Miller *et al.*, 1999).

The inventors also contemplate experiments where high-affinity antibodies raised against CeDAT should be able to partition these mutants into their proper complementation groups. Finally, since DAT plays a principle role in DA homeostasis and abnormal DA neurotransmission and has been indicated in a variety of other neuropsychiatric disorders including schizophrenia, drug addiction, Tourette's Syndrome, and Attention Deficit Hyperactivity Disorder, novel regulators identified from this screen could aid in understanding pathophysiology of these diseases (Gainetdinov *et al.*, 1998; Miller *et al.*, 1999).

Toxin Sensitivity. The inventors have shown that *C. elegans* DA neurons are vulnerable to mammalian DA neuronal toxins (Nass *et al.*, 2000). A loss of reporter expression in DA neurons was seen following 6-OHDA treatment using the GFP constructs described above. A CeDAT deficient line has been established ((Nass *et al.*, 2000; J. Duerr and J. Rand, personal communication). Using this line the inventors contemplate that CeDAT antagonist will reduce or eliminate toxin sensitivity. If the 6-OHDA action truly reflects DA neuron toxicity, one may

also expect to see behavioral deficits like those seen after laser and genetic ablations of DA neurons (Lints and Emmons, 1999; Duerr *et al.*, 1999; Sawin *et al.*, 2000).

5 The inventors also contemplate identifying molecular determinants of toxin sensitivity of DAT. For example, 6-OHDA and MPP⁺ appear to confer DA neuronal death via impairing cellular ROS homeostasis, although their mechanism of action *in vivo* remains unclear (Yahr and Bergmann, 1986; Lotharius *et al.*, 1999). Both toxins inhibit complexes in the mitochondrial transport chain (6-OHDA, complex I and IV, MPP⁺, complex I) (Glinka *et al.*, 1997; Kopin, 1992). 6-OHDA is also easily oxidized to generate hydroxyl radicals and the superoxide ion and
10 can covalently interact with cysteinyl groups to inhibit protein function (Cohen and Heikkila, 1974; Graham *et al.*, 1978). It is debatable whether MPP⁺ confers toxicity via inhibition of mitochondrial function since a cell which does not have mitochondria is also sensitive to the toxin (Khan *et al.*, 1997). The screen of the present invention is capable of identifying genes involved directly or indirectly in the increased generation of ROS (loss-of-function mutants), as
15 well as those which protect the cell from oxidation (gain-of-function mutants). Since DA itself may contribute to DA neurotoxicity (see below), mutations within the DA synthesizing (*e.g.*; TH, AAAD), DA metabolizing (*e.g.*, monoamine oxidase), and vesicle sequestration (*e.g.*, VMAT, Vacuole H⁺-ATPase [V-ATPase]) pathways are contemplated (Table 1) (Miller *et al.*, 1999). Novel targets that would normally induce neuronal degeneration, independent of
20 oxidative pathways and those involved in DA metabolism and sequestration, will also be identified by the screening methods described here.

If DA-ergic neuronal degeneration is dependent on the cell death or a necrotic pathway, then mutants within these pathways will also be identified by the present screening methods.
25 Recent investigations suggest both neurotoxins may rely on products of programmed cell death pathways (see below). 6-OHDA causes shrinkage of DAergic cells, nuclear condensation and DNA fragmentation (*i.e.*, apoptotic characteristics) *in vitro*, while MPP⁺ does not induce these events (Choi *et al.*, 1999). Morphological and biochemical correlates of MPP⁺ toxicity *in vitro* are more consistent with necrosis rather than apoptosis displaying mitochondrial swelling and
30 scattered heterochromatin (Choi *et al.*, 1999). Interestingly, the anti-apoptotic protooncogenes in the Bcl-2 class can protect against both MPP⁺ and 6-OHDA toxicity *in vitro* and *in vivo*, suggesting overlap between the two degradation pathways (Offen *et al.*, 1998; Oh, *et al.*, 1998). Apoptosis and necrotic death are both well-defined events in *C. elegans*, with apoptosis causing cytoplasmic shrinkage, mitochondria distortion, and rapid phagocytosis, while necrosis produces

vacuoles, cell swelling, and membrane whorls (Baffi *et al.*, 1999; Robertson and Thomson, 1982; Hall *et al.*, 1997). Presently, little is known about the genes involved in the necrotic pathway, and considering that necrosis can follow neuronal insult independent of activation of the apoptotic pathway, novel genes involved in necrosis will likely be identified (Roy and Sapolsky, 1999).

Additionally, the present inventors envision using the *C. elegans* model described here to study the effects of these and other neurotoxins on DA neuronal degradation in assays independent of the genetic screens.

The inventors also contemplate using the existing wealth of *C. elegans* mutants (see for example mutants listed at <http://www.elegans.swmed.edu/>). For example, PD may be the result of excess DA in the cytoplasm due to the weakening of the ion gradients required for DA sequestration in vesicles. To determine if endogenous DA levels play a role in toxin-mediated neural degeneration, the inventors contemplate crossing the reporter line P_{CeDAT}::GFP to worm lines deficient in DA synthesis such as the TH (*cat-2*) or AAAD (*bas-1*) null mutants and test for sensitivity to 6-OHDA. There is also evidence that the endogenous VMAT levels may predict the sensitivity of the DA neurons to the toxins: the lower the amount of VMAT, greater the level of DA in the cytoplasm and therefore the greater the likelihood of neural degeneration (Miller *et al.*, 1999). It is contemplated that this possibility will be tested by crossing the GFP reporter line into the VMAT null strain (*cat-1*) followed by screening for toxin induced neurodegeneration. It is also contemplated that reserpine an agent which blocks VMAT and inhibits DA vesicular storage, which causes a reduction in intracellular DA both in humans and worms), will be applied to WT worms prior to toxin exposure to determine the role of VMAT on neurotoxicity (Duerr *et al.*, 1999; Sulston *et al.*, 1975; Miller *et al.*, 1999).

Cell Death. Additionally the inventors contemplate using the *C. elegans* model to explore genes involved in cell death. *C. elegans* was the first organism in which factors regulating cell death were identified on a genetic level. Mutants in these genes are available for evaluating their roles in DA neurodegeneration (Hengartner, 1997). For example, CED-3 is a member of the ICE-family of cysteine proteases or caspases, which are known to function in both nematode and mammalian apoptosis. CED-3 activity is required for programmed cell death and a *ced-3* (lof) mutation blocks all programmed cell death in the nematode (Ellis and Horvitz, 1986). CED-4 is required for CED-3 activation. CED-9 is homologous to mammalian Bcl-2 and

negatively regulates CED-3 through interactions with CED-4. Gain-of-function mutations in *ced-9* also blocks apoptosis (Hengartner *et al.*, 1992). The present inventors contemplate crossing the P_{CeDAT}::GFP line into the *ced* mutants to directly test if apoptosis is involved in 6-OHDA or MPP⁺ neurotoxicity.

5

The inventors also contemplate that the CeDAT-directed screens will yield molecular insights into presynaptic DAT regulation and DA neurodegeneration. Due to the high similarity between the human and worm nervous systems at the molecular level, the molecules identified by these screens are envisioned to be of general relevance.

10

: All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred
15 embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be
20 achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

5

Baffi, Palkovits, Castillo, Mezey, Nikodem, "Differential expression of tyrosine hydroxylase in catecholaminergic neurons of neonatal wild-type and Nurr1-deficient mice," *Neuroscience*, 93:631-642, 1999.

10

Bargmann, "Neurobiology of the *Caenorhabditis elegans* genome," *Science*, 282:2028-2033, 1998.

BLAST search performed at WormBase using the WormPep Version 18 at <http://www.wormbase.org/perl/ace/elegans/searches/blast>

Brenner, "The genetics of *Caenorhabditis elegans*," *Genetics*, 77:71-94, 1974.

15

C. elegans Sequencing Consortium. Genome Sequence of the Nematode *C. elegans*: A platform for investigating biology, *Science*, 282:2012-2018, 1998.

Caenorhabditis elegans, www.elegans.swmed.edu/

20

Castillo, Baffi, Palkovits, Goldstein, Kopin, Witta, Magnuson, Nikodem, "Dopamine biosynthesis is selectively abolished in substantia nigra/ventral tegmental area but not in hypothalamic neurons in mice with targeted disruption of the Nurr1 gene," 11:36-46, 1998.

Chalfie, Tu, Euskirchen, Ward, Prasher, "Green fluorescent protein as a marker for gene expression," *Science*, 263:802-805, 1994.

Chen N, Reith, *Eur J Pharmacol*, Sep 29; 405(1-3):329-339, 2000.

25

Choi, Yoon, Oh, Choi, O'Malley, Oh, "Two distinct mechanisms are involved in 6-hydroxydopamine- and MPP⁺-induced dopaminergic neuronal cell death: role of caspases, ROS, and JNK," *J. Neurosci. Res.*, 57:86-94, 1999.

Cohen and Heikkila, "The generation of hydrogen peroxide, superoxide radical, and hydroxyl radical by 6-hydroxydopamine, dialuric acid, and related cytotoxic agents," *J. Biol. Chem.*, 249:2447-2452, 1974.

30

Duerr, Frisby, Gaskin, Duke, Asermely, Huddleston, Eiden, Rand, "The *cat-1* gene of *Caenorhabditis elegans* encodes a vesicular monoamine transporter required for specific monoamine-dependent behaviors," *J. Neuroscience*, 9:72-84, 1999.

Ellis and Horvitz, "Genetic control of programmed cell death in the nematode *C. elegans*," *Cell*, 44:817-829, 1986.

- Emmons, Klass, Hirsh, "Analysis of the constancy of DNA sequences during development and evolution of the nematode *Caenorhabditis elegans*," *Proc. Natl. Acad. Sci. USA* 76:1333-1337, 1979.
- 5 Fradkov, Chen, Ding, Barsova, Matz, Lukyanov, "A novel fluorescent protein from *Discosoma* coral and its mutants possesses a unique far-red fluorescence," *FEBS Lett*, 479(3):127-30, 2000.
- Gainetdinov, Jones, Fumagalli, Wightman, Caron, Re-evaluation of the role of the dopamine transporter in dopamine system homeostasis, *Brain Res. Rev.*, 26:148-153, 1998.
- GenBank accession Number AF115382.
- 10 Giros and Caron, "Molecular characterization of the dopamine transporter," *Trends Pharm. Sci.*, 14:43-49, 1993.
- Giros, El Mstikawy, Godinot, Zheng, Han, Yang-Feng, Caron, "Cloning, pharmacological characterization, and chromosome assignment of the human dopamine transporter," *Mol. Pharm.*, 42:383-390, 1992.
- 15 Giros, Jaber, Jones, Wightman, "Hyperlocomotion and indifference to cocaine and amphetamine in mice lacking the dopamine transporter," *Nature*, 379:606-612, 1996.
- Glinka, Gassen, Youdim, "Mechanism of action of 6-hydroxydopamine neurotoxicity," *J. Neural. Transm. Suppl.*, 50:55-66, 1997.
- Graham, Tiffany, Bell, Gutknecht, "Autooxidation versus covalent binding of quinones as the mechanism of toxicity of dopamine, 6-hydroxydopamine, related compounds toward
- 20 C1300 neuroblastoma cells *in vitro*," *Mol. Pharmacol.*, 14:644-653, 1978.
- Hall, Gu, Garcia-Anoveros, Gong, Chalfie, Driscoll, "Neuropathology of degenerative cell death in *Caenorhabditis elegans*," *J. Neuroscience*, 17:1033-1045, 1997.
- Hengartner, "Cell death," In : *C. elegans II*, Riddle, Blumenthal, Meyer, Priess (Eds), New York: Cold Spring Harbor Laboratory Press; 383-415, 1997.
- 25 Hengartner, Ellis, Horvitz, "*Caenorhabditis elegans* gene *ced-9* protects cells from programmed cell death," *Nature*, 356:494-499, 1992.
- Hope laboratory worm pages. URL: <http://bgypc086.leeds.ac.uk/>
- Javitch, D'Amato, Strittmatter, Snyder, "Parkinsonism-inducing neurotoxin, N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine: uptake of the metabolite N-methyl-4-pheynylpyridine by
- 30 dopamine neurons explains selective toxicity," *Proc. Nat. Acad. Sci. USA*, 82:2173-2177, 1985.

- Jayanthi, Apparsundaram, Malone, Ward, Miller, Eppler, Blakely, "The *Caenorhabditis elegans* gene T23G5.5 encodes an antidepressant- and cocaine-sensitive dopamine transporter," *Mol. Pharmacol.*, 54:601-609, 1998.
- Jenner, "Oxidative mechanisms in nigral cell death in Parkinson's disease," *Mov. Disord.*, 13:24-34, 1998.
- Johnsen and Baillie, "Mutation," In: *C. elegans II*, Riddle, Blumenthal, Meyer, Priess (Eds.), New York: Cold Spring Harbor Press; 79-95, 1997.
- Khan, Filiano, King, Przedborski, "Is Parkinson's disease (PD) an extra-mitochondrial disorder?" *Neurology*, 48:A201, 1997.
- Kitayama, Shimada, Uhl, "Parkinsonism-inducing neurotoxin MPP+: uptake and toxicity in nonneuronal COS cells expressing dopamine transporter cDNA," *Ann. Neurol.*, 32:109-111, 1992.
- Kopin, "Features of the dopaminergic neurotoxin MPTP," *Acad Sci*, 648:96-104, 1992.
- Le, Conneely, He, Jankovic, Appel, "Reduced Nurr1 expression increases the vulnerability of mesencephalic dopamine neurons to MPTP-induced injury," *J. Neuroscience*, 19(5):2218-2221, 1999.
- Lewis and Fleming, "Basic culture methods in *Caenorhabditis elegans*: modern biological analysis of an organism," *Methods Cell Biol*, 48:3-29, 1995.
- Lints and Emmons, "Genetic Analysis of Dopaminergic Neurons in the Nematode *Caenorhabditis elegans*," In: *Development of Dopaminergic Neurons*, di Porzio, Pernas-Alonso, Perrone-Capano (Eds.), Austin, TX, R.G. Landes Company, 175-190, 1999.
- Lints and Emmons, "Patterning of dopaminergic neurotransmitter identity among *Caenorhabditis elegans* ray sensory neurons by a TGF β family signaling pathway and a *Hox* gene," *Development*, 126:5819-5831, 1999.
- Liu and Sternberg, "Sensory regulation of male mating behavior in *Caenorhabditis elegans*," *Neuron*, 14:1-20, 1995.
- Loer and Kenyon, "Serotonin-deficient mutants and male mating behavior in the nematode *Caenorhabditis elegans*," *J. Neuroscience*, 13:5407-5417, 1993.
- Lotharius, Dugan, O'Malley, "Distinct mechanisms underlie neurotoxin-mediated cell death in cultured dopaminergic neurons," *J. Neuroscience*, 19:1284-1293, 1999.
- Mello and Fire, "DNA transformation in *Caenorhabditis elegans*: modern biological analysis of an organism," Epstein and Shakes, (Eds.), *Methods Cell Biol*, 48:451-482, (Chapter 19), 1995.

- Mello and Fire, "DNA transformation in *Caenorhabditis elegans*: modern biological analysis of an organism," Epstein and Shakes, (Eds.), *Academic Press*, 1995.
- Mello, Kramer, Stinchcomb, Ambros, "Efficient gene transfer in *C. elegans*: Extrachromosomal maintenance and integration of transforming sequences," *EMBO J.*, 10, 3959-3970, 1991.
- 5 Kramer, French, Park, Lohanson, "The *Caenorhabditis elegans* rol-6 gene, which interacts with the sqt-1 collagen gene to determine organismal morphology, encodes a collagen," *Mol. Cell. Biol.*, 10:2081-2090, 1990.
- Miller, Desai, Hardin, Piston, Patterson, Fleenor, Xu, Fire, "Two-color GFP expression for *C. elegans*," *Biotechniques*, 26:914-921, 1999a.
- 10 Miller, Gainetdinov, Levey, Caron, "Dopamine transporters and neuronal injury," *Trends Pharm. Sci.*, 20(10):424-429, 1999b.
- Nass, Duerr, Rand, Miller, Blakely, "6-OHDA sensitivity of dopaminergic neurons in *C. elegans*: role of the dopamine transporter and cell death pathways," *Society Neuroscience*, Abstract: 2000.
- 15 Nirenberg, Chan, Vaughan, Uhl, Kuhar, Pickely, "Immunogold localization of the dopamine transporter: an ultrastructural study of the rat ventral tegmental area," *J. Neuroscience*, 17:5255-5262, 1997.
- Offen, Beart, Cheung, Pascoe, Hochman, Gorodin, Melamed, Bernard, Bernard, "Transgenic mice expressing human Bcl-2 in their neurons are resistant to 6-hydroxydopamine and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine neurotoxicity," *Proc. Natl. Acad. USA*, 20 95:5789-5794, 1998.
- Oh, Choi, Kim, Seo, O'Malley, Oh, "Overexpression of HA-Bax but not Bcl-2 or Bcl-XL attenuates 6-hydroxydopamine-induced neuronal apoptosis," *Exp. Neurol.*, 154:193-198, 1998.
- 25 Pickel, Nirenberg, Milner, "Ultrastructural view of central catecholaminergic transmission: immunocytochemical localization of synthesizing enzymes, transporters, and receptors," *J. Neurocytology*, 25:843-856, 1996.
- Pifl, Giros, Caron, "Dopamine transporter expression confers cytotoxicity to low doses of the Parkinsonism-inducing neurotoxin 1-methyl-4-phenylpyridinium," *J. Neuroscience*, 30 13(10):4246-4253, 1993.
- Riddle, Blumenthal, Meyer, Priess, "Introduction to *C. elegans*," In : *C. elegans II*, New York: Cold Spring Harbor Laboratory Press, 1-22, 1997.
- Robertson and Thomson, "Morphology of programmed cell death in the ventral nerve cord of *Caenorhabditis elegans*," *J. Embryol. Exp. Morphol.*, 67:89-100, 1982.

- Rongo, Whitfield, Rodal, Kim, Kaplan, "LIN-10 is a shared component of the polarized protein localization pathways in neurons and epithelia," *Cell*, 94:751-759, 1998.
- Roy and Sapolsky, "Neuronal apoptosis in acute necrotic insults: why is this subject such a mess?" *Trends in Neuroscience*, 22:419-422, 1999.
- 5 Sachs and Jonsson, "Mechanism of action of 6-hydroxydopamine," *Biochem. Pharmacol*, 24:1-8, 1975.
- Sawin, Ranganathan, Horvitz, "*C. elegans* locomotory rate is modulated by the environment through a dopaminergic pathway and by experience through a serotonergic pathway," *Neuron*, 26:619-631, 2000.
- 10 Stinchcomb, Shaw, Carr, Hirsch, "Extrachromosomal DNA transformation of *Caenorhabditis elegans*, *Mol. Cell Bio.*, 5:3484-3496, 1985.
- Sulston and Horvitz, "Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*, *Dev. Biol.*, 56:110-156, 1977.
- Sulston, Dew, Brenner, "Dopaminergic neurons in the nematode *Caenorhabditis elegans*," *J. Comp. Neurol.*, 163:215-226, 1975.
- 15 Sulston, Du, Thomas, Wilson, Hillier, Staden, Halloran, Green, Thierry-Mieg, Qiu, Dear, Coulson, Craxton, Durbin, Berks, Metzstein, Hawkins, Ainscough, Waterson, "The *C. elegans* genome sequencing project: a beginning," *Nature*, 356:37-41, 1992.
- Sulston and Hodgkin, "Methods," In: *The nematode Caenorhabditis elegans*, Wood and the Community of *C. elegans* Researchers (eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 587-606, 1988.
- 20 White, Southgate, Thompson, Brenner, "The structure of the nervous system of the nematode *Caenorhabditis elegans*," *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 314:1-340, 1986.
- Wood, "Introduction to *C. elegans* Biology," In: *The nematode C. elegans*, Wood (ed.), New York: Cold Spring Harbor Laboratory Press; 1-16, 1988.
- 25 Yahr and Bergmann, In: *Advances in Neurology: Parkinson's Disease*, New York: Raven Press, 1986.

CLAIMS

1. A recombinant *C. elegans* that expresses a detectable marker in a dopamine neuron.
- 5 2. The recombinant *C. elegans* of claim 1, wherein the detectable marker is further defined as a marker that can be visually detected.
3. The recombinant *C. elegans* of claim 1, wherein the detectable marker is further defined as a marker that can be spectroscopically detected.
- 10 4. The recombinant *C. elegans* of claim 1, wherein the detectable marker is a green fluorescent protein.
5. The recombinant *C. elegans* of claim 1, wherein the detectable marker is a yellow
15 fluorescent protein.
6. The recombinant *C. elegans* of claim 1, wherein the detectable marker is a blue fluorescent protein.
- 20 7. The recombinant *C. elegans* of claim 1, wherein the detectable marker is a red fluorescent protein.
8. The recombinant *C. elegans* of claim 1, wherein the detectable marker is β -galactosidase.
- 25 9. The recombinant *C. elegans* of claim 1, wherein the detectable marker is under the control of a promoter.
10. The recombinant *C. elegans* of claim 9, wherein the detectable marker is an antigenic polypeptide.
- 30 11. The recombinant *C. elegans* of claim 9, wherein said promoter is a tissue-specific promoter.

12. The recombinant *C. elegans* of claim 11, wherein the tissue-specific promoter is a neuronal promoter.
13. The recombinant *C. elegans* of claim 12, wherein the neuronal promoter is a dopamine transporter promoter.
14. The recombinant *C. elegans* of claim 13, wherein the neuronal promoter is a tyrosine hydroxylase promoter.
15. A method of screening for substances that affect neuronal viability comprising:
- a) providing a recombinant *C. elegans* that expresses a detectable marker in a neuronal cell;
 - b) exposing said *C. elegans* to a candidate substance; and
 - c) detecting a change in the expression of the marker relative to the expression of the marker before said exposing;
- wherein a change in the expression of the marker corresponds to a change in the viability of the neuron.
16. The method of claim 15, further comprising detecting the expression of the marker in the neuronal cell in the absence of said candidate substance.
17. The method of claim 15, wherein said substance is a neurotoxic substance.
18. The method of claim 17, wherein the neurotoxic substance is 6-hydroxydopamine, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, or 5,7-di-hydroxy tryptamine.
19. The method of claim 17, wherein the neurotoxic substance is 6-hydroxydopamine.
20. The method of claim 17, wherein the neurotoxic substance is a generator of free radical species.
21. The method of claim 15, wherein said substance is a neuroprotective substance.

22. The method of claim 21, wherein the neuroprotective substance is a dopamine transporter antagonist.
23. The method of claim 22, wherein the dopamine transporter antagonist is imipramine.
- 5 24. The method of claim 22, wherein the dopamine transporter antagonist is an amphetamine.
25. The method of claim 22, wherein the dopamine transporter antagonist is cocaine.
- 10 26. The method of claim 21, wherein the neuroprotective substance is a free radical scavenger.
27. The method of claim 15, further comprising the step of exposing said *C. elegans* to a known neurotoxin prior to step b).
- 15 28. The method of claim 15, further comprising the step of exposing said *C. elegans* to a known neurotoxin after step b).
29. The method of claim 15, wherein the substance is encoded by a polynucleotide.
- 20 30. The method of claim 15, wherein the polynucleotide encodes a dopamine transporter regulatory gene, or a gene that suppresses free radical generation.
31. The method of claim 15, wherein the substance is a polypeptide.
- 25 32. The method of claim 31, wherein the polypeptide encodes a dopamine transporter regulatory polypeptide, or a polypeptide that suppresses free radical generation.
33. The method of claim 15, wherein the substance is a naturally occurring product.
- 30 34. The method of claim 15, wherein the substance is a man-made chemical.
35. The method of claim 34, wherein the man-made chemical is a monoamine oxidase inhibitor.

36. The method of claim 35, wherein the monoamine oxidase inhibitor is a hydrazine derivative.
- 5 37. The method of claim 36, wherein the hydrazine derivative is phenelzine or isocarboxazid.
38. The method of claim 35, wherein the monoamine oxidase inhibitor is a non-hydrazine derivative.
- 10 39. The method of claim 38, wherein the non-hydrazine derivative is tranylcypromine, or pargyline.
40. The method of claim 15, wherein the substance is an environmental toxin.
- 15 41. The method of claim 40, wherein the environmental toxin is a pesticide, or a herbicide.
42. The method of claim 41, wherein the pesticide is rotenone.
43. The method of claim 15, further comprising:
- 20 a) exposing said *C. elegans* to a known neurotoxin; and
b) detecting a change in expression of said marker.
44. The method of claim 15, wherein the change in marker expression can be an increase in the marker.
- 25 45. The method of claim 15, wherein the change in marker expression can be a decrease in the marker.
46. The method of claim 15, wherein the detectable marker is further defined as a marker that
- 30 can be visually detected.
47. The method of claim 15, wherein the detectable marker is further defined as a marker that can be spectroscopically detected.

48. The method of claim 47, wherein the detectable marker is a green fluorescent protein.
49. The method of claim 47, wherein the detectable marker is a yellow fluorescent protein.
- 5 50. The method of claim 47, wherein the detectable marker is a blue fluorescent protein.
51. The method of claim 47, wherein the detectable marker is a red fluorescent protein.
52. The method of claim 15, wherein the detectable marker is β -galactosidase.
- 10 53. The method of claim 15, wherein the detectable marker is under the control of a promoter.
54. The method of claim 15, wherein the detectable marker is an antigenic polypeptide.
- 15 55. The method of claim 15, wherein the detectable marker is under the control of a neuronal-specific promoter.
56. The method of claim 55, wherein the neuronal-specific promoter is a dopamine transporter promoter.
- 20 57. The method of claim 55, wherein the neuronal-specific promoter is a tyrosine hydroxylase promoter, a *cha-1* promoter, an *acr-2* promoter, an *unc-30* promoter, an *unc-4* promoter, or an *asi* promoter.
- 25 58. The method of claim 15, wherein the neuronal cell comprises a dopaminergic neuron.
59. The method of claim 15, wherein the neuronal cell comprises a cholinergic neuron.
- 30 60. The method of claim 15, wherein the neuronal cell comprises a GABA-ergic neuron.
61. The method of claim 15, wherein the neuronal cell comprises a glycinergic neuron.
62. The method of claim 15, wherein the neuronal cell comprises a serotonergic neuron.

63. The method of claim 15, wherein the neuronal cell comprises a glutamatergic neuron.
64. The method of claim 15, wherein the neuronal cell comprises a peptidergic neuron.
- 5 65. A method of screening for substances that can inhibit neuronal cell death comprising:
- a) providing a recombinant *C. elegans* that expresses a detectable marker in a neuronal cell;
 - b) exposing said *C. elegans* to a known neurotoxin and a candidate substance;
 - 10 c) detecting expression of the marker; and
 - d) comparing the expression of the marker to the expression of the marker in the absence of the candidate substance.
66. The method of claim 65, wherein the *C. elegans* is exposed to the neurotoxin prior to the candidate substance.
- 15 67. The method of claim 65, wherein the *C. elegans* is exposed to the candidate substance prior to the neurotoxin.
68. The method of claim 65, wherein the detectable marker is a green fluorescent protein.
- 20 69. The method of claim 65, wherein the detectable marker is a yellow fluorescent protein.
70. The method of claim 65, wherein the detectable marker is a blue fluorescent protein.
- 25 71. The method of claim 65, wherein the detectable marker is a red fluorescent protein.
72. The method of claim 65, wherein the detectable marker is β -galactosidase.
- 30 73. The method of claim 65, wherein the detectable marker is under the control of a promoter.
74. The method of claim 65, wherein the detectable marker is an antigenic polypeptide.

75. The method of claim 65, wherein the detectable marker is under the control of a neuronal-specific promoter.

76. The method of claim 75, wherein the neuronal-specific promoter is a dopamine transporter promoter.

77. The method of claim 75, wherein the neuronal-specific promoter is a tyrosine hydroxylase promoter, a *cha-1* promoter, an *acr-2* promoter, an *unc-30* promoter, an *unc-4* promoter, or an *asi* promoter.

10

78. A method of screening candidate substances to identify a substance that can be used for prevention and/or therapy of neurodegenerative diseases comprising:

- a) obtaining a recombinant *C. elegans* that expresses a detectable marker in a neuronal cell under the control of a neuronal-specific promoter;
- 15 b) exposing said *C. elegans* to a known neurotoxin and a candidate substance;
- c) detecting expression of the marker; and
- d) comparing the expression of the marker to the expression of the marker in the absence of the candidate substance.

20 79. The method of claim 78, wherein the *C. elegans* is exposed to the neurotoxin prior to the candidate substance.

80. The method of claim 78, wherein the *C. elegans* is exposed to the candidate substance prior to the neurotoxin.

25

81. The method of claim 78, wherein the neurodegenerative disease is selected from Parkinson's disease, Alzheimer's disease, Huntington's disease, a transmissible spongiform encephalopathy (TSE), a familial amyloid polyneuropathy (FAP), a prion diseases, a Tauopathy, a Trinucleotide disease, amyotrophic lateral sclerosis (ALS) or multiple system atrophy.

30

82. The method of claim 78, wherein the detectable marker is a green fluorescent protein.

83. The method of claim 78, wherein the detectable marker is a yellow fluorescent protein.

84. The method of claim 78, wherein the detectable marker is a blue fluorescent protein.
85. The method of claim 78, wherein the detectable marker is a red fluorescent protein.
- 5 86. The method of claim 78, wherein the detectable marker is β -galactosidase.
87. The method of claim 78, wherein the detectable marker is under the control of a promoter.
- 10 88. The method of claim 78, wherein the detectable marker is an antigenic polypeptide.
89. The method of claim 78, wherein the detectable marker is under the control of a neuronal-specific promoter.
- 15 90. The method of claim 78, wherein the neuronal-specific promoter is a dopamine transporter promoter.
91. The method of claim 78, wherein the neuronal-specific promoter is a tyrosine hydroxylase promoter, a *cha-1* promoter, an *acr-2* promoter, an *unc-30* promoter, an *unc-4* promoter, or an *asi* promoter.
- 20 92. A method of screening for substances that modulate dopamine transporter function comprising:
- 25 a) obtaining a recombinant *C. elegans* that expresses a detectable marker in a dopaminergic neuronal cell;
- b) exposing said *C. elegans* to a candidate substance;
- c) exposing said *C. elegans* to a neurotoxin that requires a dopamine transporter for intracellular access; and
- 30 d) detecting any change in the expression of the GFP after step c).
93. The method of claim 92, wherein the candidate substance blocks transport by the dopamine transporter.

94. The method of claim 92, wherein the candidate substance increases transport by the dopamine transporter.
95. The method of claim 92, wherein the neurotoxin is a addictive substance.
- 5 96. The method of claim 92, wherein the addictive substance is selected from cocaine, amphetamines, methyl-phenidate.
- 10 97. The method of claim 92, used to identify substances that provide therapy for neurological diseases involving dopamine transporter function.
- 15 98. The method of claim 97, wherein the neurological diseases involving dopamine transporter function are Schizophrenia, addiction disorders, attention deficit hyperactivity disorder, psychoses, Tourette's syndrome, or Parkinson's disease.
199. A method of screening for molecules that modulate neuronal signaling comprising:
- a) obtaining a recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a knockout and/or a mutant for a component of neuronal signaling;
 - 20 b) obtaining a second recombinant *C. elegans* that expresses a detectable protein in a neuronal cell which is a or a second mutant for a component of neuronal signaling;
 - c) comparing the differences in neuronal viability when exposed to a neurotoxic substance in the *C. elegans* of step a) with the *C. elegans* of step b); and
 - 25 d) identifying the genetic component of the mutation.
100. The method of claim 99, further comprising isolating the genetic component.




FIG. 1A




FIG. 1B

2/8

GENETIC SCREEN FOR REGULATORS OF CeDAT FUNCTION,
REGULATION, AND DA NEURON TOXIN SENSITIVITY




FIG.2

3/8




FIG.3A




FIG.3B




FIG.3C




FIG.3D




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D

5/8




FIG. 4F




FIG. 4E




FIG. 5B




FIG. 5D




FIG. 5A




FIG. 5C

7/8




FIG.5E

8/8




FIG. 6B




FIG. 6D




FIG. 6A




FIG. 6C

SEQUENCE LISTING

<110> VANDERBILT UNIVERSITY
BLAKELY, RANDY D.
NASS, RICHARD
MILLER, DAVID M.

<120> ASSAY FOR TOXIN INDUCED NEURONAL DEGENERATION AND
VIABILITY IN C. ELEGANS

<130> VBLT:007-WO

<140> UNKNOWN

<141> 2002-06-18

<150> 09/888,233

<151> 2001-06-22

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 719

<212> DNA

<213> C. ELEGANS

<400> 1

```
ccatgaaatg gaacttgaat ccagttttca ctaaaacgac ctcatacact ttctctcgta 60
tcctcaaaat atctatgaca ttatcattag cttecgtagt ttcatttctt tcaaataatta 120
tgcattctta aattccgata cccgcgtgca aaagtgtctt attgagcaac tttgggatca 180
tatgtacaca ccaatgccct ttcccaaat cttttcctgt ccttttctct aaaaacaata 240
aatccatgcc tattccagta tgacccttt gaagcagata taatcgaca aacatatata 300
catagctcgg ataaatgtag aaaaagaaga aaagaagtat aagtagatag atgctttccg 360
gcaattatcc accgcaccgt agtcctcacc aactgagact gcgtcgtag gagacgcga 420
catgattcag aagcagaatt tggaagaaaa acgacgatga tattgaggct ggcacacata 480
caccggaata ttgcacatgc caccacatct agattccaag gcaatctcta cctcttccca 540
ttctttcggg tttttgttgc tgacaagaaa agtggatagc tacgggctca atgagctgat 600
tttattttta aatatcttaa aactatacta gattcatgtg ttttcaggct catattccaa 660
attagtcgaa aagctgatcc cgctacggtt tactcgaatc tcaacaattt ttagccatg 719
```

<210> 2

<211> 10851

<212> DNA

<213> Homo sapiens

<400> 2

```
aaaggcgaga ctgaatatgt ttaatagaga catgacagat attttttaaa agacacgaat 60
ccaaattcta gagatgaaaa ccacaatgtc tcagatgcaa aatgcactgg ttgagaggag 120
gggtggattc aacatctcag aagaaaagga tacatgaact tgaagacata gatggcagtg 180
caaactgttc aaaatgaaat aaagaaaaaa gacaaaaaaa tgaaccaagc atcagagaac 240
tgtgggactt caagccacct aatatgcaag ccataggaat cctcaaaaca gatcatgtct 300
aaacaatgtc attttatgat agcatcagtt caccaaaagg acaaattgtt atgtacctaa 360
taacatagct tcataatgaa ttccggcaaaa actgatagaa ctgcaaggag acatagagca 420
gttcacacaa ttatacttgg agatttcaat acctctttcc caatagttga tagaacagat 480
acacagaaac ttttaaggat acagaagact tcaaaaacag tacgaactaa catgacctat 540
ttgatattta tacaataatt catccaataa tagcagagtg cacattcttt tcaaatgcat 600
```

```

gtgaaaagct taccaagatc gaccacagtc cagaccataa aattcccttt agggattcag 660
gtcatgtataa gcaagttctc caacaaagag gcaaataaat ttgatacca aatgaaaaat 720
acatttggca aaaaataaag aagcatatcc agacaatctt ctaatatctg gaaactaatc 780
acatatctaa ataacaatgg gtcaaagagg acatcaaaag ggaaatcaga aaatattttg 840
aactgaatga aaacaaaaca acatatcaaa ccttttggaa tttggctaaa tcagaactta 900
gaagaaaatc tagacatgaa gcctgcctca gaaaggaata atgatgaaaa attgatggca 960
gcctccaccc gaagaaatcc catgcgggac tttggaagaa attgaaaagt tgattccaaa 1020
attcatatga aaaagcacag gaccgagaat atgcaaaaaca actttgaaaa tgaacaaagt 1080
tgtctgactt acattactct atttaaaaat gtattacaaa aggaccataa tgaagatagt 1140
gccattttta tctcctcagg ctgccacaac aaaatacacc agatttgggg gcttaacagc 1200
agacatttgt ttctcacagt tctggagact ggaaatccaa catcaaggca gctgattcag 1260
ttcctgggtg ggcctctctt cctggcttgc agatggccgc cttcttactc tatcttcaca 1320
tggcagaaaag taaaatggag agagcttcct cttcttataa ggccacagtc ctctgggtt 1380
atgaccccac ccttaacaatt taatttaacc taatcacctc ctaaagtccc tatcctccag 1440
ataaagtcac ctgtcggggg ttatggcttt aacctatgaa tttgggggta aacaattcag 1500
tccacagcaa gtgtgtgtatt agcatcaaaa tggatttaaca gatcataaaa caggagagccc 1560
agaaataaac ccacacacat acatacaact gacttttcaa caaagggtgca aaagcaactc 1620
agcagaacac tgaacaactg gctcttcgga ggtaaagtgg tgaacttcaa ttgggtgtctt 1680
acaccaagct agtttaaaaa tgcattctat gtgtaaatat gcaatctata attttaaaaa 1740
gttttagagga aaacttagga gaaaatcttg atgatcatga atttggtaaa gatttcttaa 1800
atacaacact aaaagcagga tttgtgaaag aaaaatggat aaattcgact tcatcaaaaat 1860
taataactct tctctgaaag aatctattaa gagaatgaaa agacagtcca agactgggaa 1920
aaatatttgc aaagtaccca tctgataagg gactgtgtac cagaatagta aagaactctc 1980
aaagctcaat taaacaaccc aaaggtgcaa aagatttgaa cagacactta acacgaagtg 2040
gcactgatgc agctaacaca tgagaaattg ttcaaagatc acgctcatta gagcagtata 2100
ggagatatta ctgtaaacat attaggtatg ctaaaattta aaagactgac cataactaag 2160
attcattaga atatgaagta actaaaatta tcatacactg ttcatggaaa tataaaatga 2220
tacaactact ttgcaaaaaga atttaacagt ttttaaaaaa atagttaaac atatatctta 2280
ccaaatgacc cagccattcc acaactaagt acttcttacc taccctaaag taatgagagc 2340
ctatgttcaa agacttgcac acaaatgttc atagtagctt tatttgtaac agccagatgt 2400
ccatcaatag gcagatggaa aagccaagta catcatatcc catatccaca caatggatac 2460
tacaataaac taaaagggaa ttaactgccg atgcaatgat gtggattctt ctcaaataatg 2520
ctgaatgcaa aagaacagac ccagaaaaca gtacttactg tataattcta tttatataaa 2580
attctattaa aaacacattg ttctgtaatc atagaaaaca aatcattcca ccattttggg 2640
aggccaaggg gagcttatca cttgagtttg ggagttcaag accagcctgg ccaacatagt 2700
gaaaccctgt ttctactaaa aatacaaaaa tttagctgggt gtgggtggcg gcactctgta 2760
tcccagctac ttgggaggct gaggcaggag aattgcttga acttgaggag gcaaaggctg 2820
cagtgaagtg agatcgtgcc actgcactcc agcctagggt acagagttag actccgtctc 2880
acaaaaagaa aaaaaaaaag agcaaatcag tggttgcctg gggttgggaa gaggtaaaga 2940
gacacagaca gagggagatc agcgagggag gagcacccgc atcatcttct gtgtgatcgc 3000
gttgccatga ccccatcagt cactcaccaa atcgtacact tggaaatattg catatccatt 3060
acacctcagc aaggctgttt ttaaaaagac acaaatgccg agataattac agaaaaaaag 3120
tccaattagt tttgttagga tgggtggaatc atgaaaggca tctcctaggt cctccaatgt 3180
tgttcttttt caagattctt ttggctactc taggtcacc acatctccaa acacatttta 3240
gaaccagctg gtttaattct tcaaaaaagc aagttagaat tttgattgag attgcatgga 3300
atctgtatgt caatttatga agaactgaca tctcattaat attgagcctt tcagtcataa 3360
acatacacac tctctgtctt gtctagactt caaaagtctc tcacagcaat ggattctagg 3420
ttttcggtgc acaagtctcg gatacgttta tccccagca ttttgttctg gatgctattg 3480
taagtgaat tatttttagt ttatttttag aattgtttgc tgccagtaca tagaaataca 3540
attggttttc atcagatccc atggccttgc taaattcatt tataaattct agtagttttt 3600
tttgtaaatt cctcaagatt ttctacaac atgaaatcta ccaagaaaaa tcttgtcctc 3660
aatatttcaa tttttttgaa gcctcctgct ctgtagccag accattcgcc actgcctatg 3720
actacattta gggccatcct ctggccatcc ctccttggct agatgatgaa cctgatgcat 3780
ggtttgataa tgtgccact ttttactgg gggagctt tctttcacag aaggccatga 3840
tgacacagcc ttctactttc tgggtggggc agtgtgtgt gtatagacac atgtaatcca 3900
ggccccaacg catgcatttt ctccgttcaa tgtcataaga aactctcatg tagccagagg 3960
tctgggttaa ggaagcggtc tccatgcagc aagaatgggt gcttgttcat ggaacttacc 4020
gcgcttccag acttgtctca gccttgtgtc gtaaaccatt cactgaggac ggcaggatgc 4080
tgtatacaca acctctgtgg gtccgataac aaatagcttc tgetggggcg ctcagcttct 4140
atggtaacag agctccctga gtttctgcta tggctcaggg ctgctcttac tcaaaaggag 4200

```

aggacgtatt ttcaggagag cgttcagctc ttaattcaac tgttattttc tgcgctgcct 4260
 acaacatctc cgtctgccac agaccctcca attcccactg gttctgccag atcagtgttc 4320
 tcaaagtgtg gttctggggc cagcagtgctc agtgtcacct gagaactttc tagaaacgca 4380
 ggtcctcagg tcccacctca gacctactga acctgcgtct ctaaggatgg agcgaggtga 4440
 tctgtgcctc ctatgcatgt aaaggtttat gtaggtgact gtgatgccag ctaaagtctg 4500
 agaacaacac cagatttgtgt ggcctcctgg ggcagcagtt tgcccatcca gaccagcagc 4560
 agaatctttc ttactctggg ccctactcaa aaccaaagc tttatgaaga acacaaaaca 4620
 ggccagagaa gcatccccga gtgtcttgca cctcaccttg caaatcgga gagcctcacc 4680
 aggtatgata cttctttctt agtggtagtc ttgcttgcca gagacagcgt gttttctttc 4740
 attttggata tataatgtga ccaatcagac caatggaccc acagaaagtt caccaaggtg 4800
 gcaggttttg aaaatgtgtt attttctctt tcaattttta gcaagcacgt gtcttaccac 4860
 ggcaccttgg catttttcat tttttgctcc ccaggttgat tcagcctgat gtcacagtg 4920
 tggcagacca ctgtgacatg ctgtgcagtg gcaagacat caagatgtct gtgactaaaa 4980
 ccagcagtg ggcggaaaat gatacagctc tgagagaaga cattggaagt gtactgctgt 5040
 ccctgccatg cgagcgaaa ctatcttctt cctgaggtgg ccccgggga gggcacagg 5100
 gtgcagcagt gagcagggcc tctgcctgg agccagcgcc ttccatcatt ctcaaacct 5160
 ggacagacaa ggtctgggat ctgacctccc acacccctgg acaaaagcca ctctgccta 5220
 ctggtccctt cacccttgcc ctcccaagc cccctacccc cagggccttc ccagcaccag 5280
 tgagagtcag ggtctttaaa gtctgagctg ggctccctgg gacttctctt gaaagcacac 5340
 aggacacctt caatatagta aatacgcagg caggtaaatg actgtcatct cactgccact 5400
 gtcttctgct cctctcagat acagcactca tggggggagc atccacgctg tcttccaaat 5460
 caggcgagac acggcgcaac atccagcact gagggcctcc caggaggcag caccacagga 5520
 gggggggacc cgtggggcaa ggttgctttg gagacagcag tcagtggcca ggggctcctt 5580
 gtgggctctg cagctgcgtt cccagccagt ggggagaggt gccgagcatg ggcaggaagt 5640
 gcagaggcag gggggctcca cctgcctgca cccaacgccc tgagcccaca gcagccatag 5700
 cagcaaccac aatgataata aagccgactt ggcatTTagg gcaaagttcc aagcatgcaa 5760
 aggtcggccg tttgatcagg tctgatcagc tcataaccac actgcttcta cctgcacagt 5820
 tcacggagca ttctgttgtt gggaggtagg agaccatgg gtctggcagc tgcgctttct 5880
 ctgtgtcatc catgagccca actccgcag ttagtTTgtt cttagagcac ccaaagctcc 5940
 tttatcctaa ttcattgtgt tggagtcgg ggttgaggca ggggtggagg aatgctcttt 6000
 gtcttggcag agtgacaggt acatgcgtgt gatcactcag tggccctcc tgagtgtggc 6060
 aggtgcattc tctgtgtgct actggtcagt aaggatgtgg ctgcctggat ctgtgtgacc 6120
 tctagtccct gcaccttctt gcctgtacct tgttagcttt gggtcacaat tctgcgctcc 6180
 tgcagcgtt gcaatccct cccaaacgct gtttgctgt gtgtgtttt gtttcgagac 6240
 aggggtctcat cctgtggccc atggcgcaat ctacgtcac tacaacctcc acctccagc 6300
 ttcaagctat tctccacct ctgcctcccc agtagctggg actacaggtg tggggccacca 6360
 tgcttggtca atttttttt tcaaagtcag ggttttgcca tgtggcccag gctggtctca 6420
 aactcctggc ctcaagagat cctccttctt cggcctccca gagtgtctgg attacaagcg 6480
 tgagccctca ctctggcct gtgtattttt aatatactg aacatccatt ctctctgtgt 6540
 gttttattta acagcctccc ttagtcacct gcaaagctt ttcttggga gactgtttcc 6600
 tcaaccctgc tgctctgggg ccaagcctg gctcactct ttttattgaa acctgtgcca 6660
 tggagataat aggggttagag agatcccttc tgtggcagcc actgacacac tacagcttcg 6720
 aggtggcaca tccccctctc ctgaagtcct ctcacctccc tggcgatgaa gtcccacccc 6780
 tgatgggagg tgggtgtcagg aggccttcag gtggtcaggc caggagggct ccacctgag 6840
 gaatgggacc agtgccctca taaaacagac cccggagagc tctccccagc ccctagcgtg 6900
 gggagataca gggagagaac tgtctgcaac cccgaagcgg ccctcaccag acacagagtc 6960
 ggccaggcct tggcctcggg acaccggaac cgttagaact gaaggcttct gtgtgagccc 7020
 ccaggctgtg gagttttttg tcatggcagc ccagggggt cactaggctc ccacttgatt 7080
 ccaactcagc gtgaagtcac agccctgagt gccttctgcc tgggtgccag ccccgaggcc 7140
 ggggagcggg ggagcggggg gcggggaggg gagtgtgtgt gtgcggggag tgcggggcgg 7200
 gcgcaggggg tggggcaccg cgtgcggggc gggtagtcgc gagttaggca ccaagggtcc 7260
 ctgcctccct cactgctgag cgcgggctgc aggtggaat ggctggagag ccccgaggct 7320
 cgctgggag cccagggcag ggtgctcac ggagcatcga gggtagacgg ggaggaacgc 7380
 cggggttcgg gcgacccatg gggcgacgca cagagctggg cgcggccact cactoggtg 7440
 ccttctaagg acctggacat cctgggcctt ggcgccctgg gggtccatt cctccgcg 7500
 ctgaatggaa gaaatccgc ccgggcatct cggaaggaaa gcctcgagc ccattoggca 7560
 ctggagccgg ataccaccc cagggttttc caggcccgct ccgggaaatg ttttcttagg 7620
 cgagtgcgag gcgggcccct cggttccgat gcaggcgcaa tagatgcgg caaggcgggg 7680
 ataggctagg ggacctcggt cgcctcgagg tcgcgagac ccaaggcca cggaaggacc 7740
 cgcgtctccg cagcccgcac gccgggaagc gtgcagagtc ctcgcgggg tcccagacc 7800

gctggtcaga	gcgtggagcg	gcgggggtggg	agggacgtgg	tccccagagc	gcggggccac	7860
cgtagggggc	cctgatgggg	agggagggaa	gggtcggccc	gacgggggtcc	cagcagttcc	7920
ccgcgcgcag	ccgtcgggct	ccctccccgt	ccagctggga	gccgccagcc	ctgggcgtcc	7980
gaagatagcg	ggtgccccgg	gcagccccc	ggggtgctgg	cgagggcgca	gggcccggcc	8040
agacagttcc	cgctggaaag	gcgcccgtct	agatccgcga	cgtctcggac	ccccaggccc	8100
ccgcacccc	tgtccgaggc	tccgggacgc	gcaggacagt	ggagccgtgg	ccgcgcgttg	8160
ctcccagcca	tctgcgtccg	ggaagcgggg	gcggggggcg	ggcccgggga	ggtgaggagg	8220
aggagccagg	acgcgagggc	gaccccgtcc	gcgggagggc	ggggcggggc	ggaccctgtc	8280
tactggataa	gacccgaggc	cgaagctgag	accgccagc	gctgcggagc	gggaggggag	8340
cttcgcggaa	cgctctcggc	gccaggactc	gcgtgcaaa	cccaggcccg	ggcgccaggt	8400
tgaggccagc	gtcgctcgcg	gcatcggggc	gccccgctcc	ttccgcagac	cccgaagtgg	8460
ggcgccaggg	cggggggccg	gggcccggca	cagtctgggg	tccccgcgtc	ccgcagaccg	8520
cgcgctctcc	aaagtgcgca	acagtcgcgg	gtgcccagcg	ccccccgata	gcgccacatg	8580
ggaccctgag	gcgctccgag	gcgcgaggag	ggtgcagggc	tgccccctgg	cccgtctcaa	8640
gctcagaacc	gggtgggcac	ctggtgcagt	caccggctta	agggacgcgt	gggtgtctat	8700
ggctgtgact	cgggggtctg	gtttcttctc	gtggaattaa	cctactaagg	gtgcggcgca	8760
tcccagatcc	gatcggaatg	ggttttgtac	accgccgctc	catctcgccg	gggctttgtc	8820
tgtgttgagg	gtggtggcgg	gcgcgcgctg	cgcgctgggt	ctctggggca	ggcggggaag	8880
ccgggagagg	actcgccagg	cagcgcgcgt	tcttgttctg	ggcgcggtga	ggaaggacgc	8940
tttctaaccg	gccacatttt	gctgtgtaga	ccaaaatcgc	ctctgaggcc	ccgcgttcag	9000
gagcggggtc	aggtggcccc	agggcgccgg	cggcttgccg	gaaactcgcg	agctccgcac	9060
ccgacgccct	ctcccaacgc	ggcctcctgc	tcgcgcgcgc	gaaccccttc	gtcgggtgtt	9120
ttaccacacc	gaggggtcgt	gcgggttgag	gttgtcaccg	ggtgcgtggc	atagctcgtg	9180
atagctcatg	ggtgaggttt	tgtgcaaa	tggatgcagg	gaaagtggcc	tgttagagcc	9240
tccacctgcg	acctgcttca	gtcgttgtgt	gtgtgtgcgc	acctgtgtga	gtgtgagtgt	9300
gtatgtgtgt	aagtgtatgt	gctcgcctgt	gtgtgtgtga	gtgtgtatgt	gtgtttgtga	9360
gtctgtgtgt	gtctgtgtgt	ctgtgtgtgc	gtgcgctcga	ctgaaacacg	ctgctgctgg	9420
atccaaatga	cagaagtcgc	cctggctggg	gcgggtgtaga	cgctcctgct	ctcctgctca	9480
gcgttgccag	gggggttatg	taccgtttgg	acangatttc	ccgggttacc	ctgctggccc	9540
aagaactaat	tcccgcngang	aaacctgtc	catcctccgc	ccaactctct	cacgcggggg	9600
ggtgccacct	gccctaagtg	gatgtggctt	gtacanacac	tttttgagga	agcagttgtg	9660
atgggttatgt	ctaaactttc	tttaacagtg	gctgattttg	ctttatataa	attttgttct	9720
ttattaactg	agtataaaca	atacaagccc	aggcttggtg	gctcatgcct	gtcatctcag	9780
cactttggga	ggctgaggca	ggagatcgcc	ttgagaccag	gagttcaaaa	ccagccttgg	9840
caacaatagt	cagaccctgt	ctctacaaga	aaacaacaac	aacaacaaaa	aaacacacac	9900
aaaaataact	tagccggtgc	tgtggtgcac	acctgtagtc	tcagctgctc	aggaggctga	9960
ggtgcaagga	tcacttgaac	ctaggagggt	gaggcagtg	gttgtaatca	caactgtatt	10020
ccatcctggg	tgacagagcg	agaccctcat	ttaaaaaaga	aaaaaaaag	aaaaaagaat	10080
acagatgaac	agtcatgaag	acattattga	atgctcctag	aagattgtaa	aattgctctc	10140
tggaagtgtg	ggggaagggt	gaagtgat	ccatgcattg	ttagtagaaa	gccacgctag	10200
agctcacaca	gccttgcact	ttgataggag	tggggagggg	tgagggggaa	ggaggagcaa	10260
accagagtgt	ctgtcttgag	gcctccatgg	gccagtgcgc	cagccctgtg	gtgagggctg	10320
gcattcccag	ctcccgtgcc	ccagctgtac	catctccagg	cgtgagaagc	acccatcctt	10380
tcccagagga	atgcccgtga	atgcttcggg	gtctgccatc	cgcaacaggt	atgtccctag	10440
ccctggctga	tgaattgttg	cgttcctggt	gtgtgtttat	ttttcatatt	ggctgaagac	10500
caagagggaa	gaagcacaga	attctcaact	cccagtgctg	ccatgagtaa	gagcaaatgc	10560
tccgtgggac	tcattgtctc	cgtgggtggc	ccggctaagg	agcccaatgc	cgtggggccc	10620
aaggaggtgg	agctcatcct	tgtcaaggag	cagaacggag	tgcaattcac	cagttccacc	10680
ttaccaacc	cgcgccagag	ccccgtggag	gcccaggatc	gggagacctg	gggcaagaag	10740
atcgatttct	cctgtccgct	attggctttg	ctgtggacct	ggccaacgct	tggcggttcc	10800
cctacctgtg	ctacaaaaat	ggtggcggtg	atcccatctc	agcttccttg	a	10851

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/19230

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : G01N 33/00; A01K 67/00, 67/033; C12N 5/00, 5/02, 5/06, 5/10 US CL : 800/3, 13; 435/325, 348 According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 800/3, 13; 435/325, 348 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST, MEDLINE, BIOSIS, EMBASE, SCISEARCH, CAPLUS														
C. DOCUMENTS CONSIDERED TO BE RELEVANT														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim												
A	NASS et al. Neurotoxin-induced degeneration of dopamine neurons in <i>Caenorhabditis elegans</i> . PNAS. 05 March 2002, Vol. 99, No. 5, pages 3264-3269, see entire document.	1-100												
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents</td><td>"T" later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principles or theory underlying the invention</td></tr><tr><td>"A" document defining the general state of the art which is not considered to be of particular relevance</td><td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>"E" earlier document published on or after the international filing date</td><td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>"G" document member of the same patent family</td></tr><tr><td>"O" document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>"P" document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents	"T" later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principles or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G" document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means		"P" document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents	"T" later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principles or theory underlying the invention													
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G" document member of the same patent family													
"O" document referring to an oral disclosure, use, exhibition or other means														
"P" document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 26 AUGUST 2002		Date of mailing of the international search report 30 SEP 2002												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-9230		Authorized officer PETER PARAS JR Telephone No. (703) 308-8340 <i>Janece F...</i>												

線虫の快と不快：感覚シグナルの受容，統合， 学習・記憶の研究から見えること

毛利亮子*, 森 郁恵*

抄 録 線虫 *C. elegans* は、単純な神経系によって、多様な快、不快に応答し、それらに対して学習・記憶能力を持つ。さらに、その全細胞系譜、全神経回路網、全塩基配列がすでに決定済みであることから、神経系の発生・機能に代表される高次生命現象を個体から遺伝子レベルまで一貫して研究するための最適なモデル生物である。これらの利点を生かして、温度走性を制御する神経回路モデルが提唱され、温度走性に関与する遺伝子も次々とクローニングされてきており、哺乳類をはじめとする高等動物では非常に困難だと予想される任意の現象を、神経回路レベルで包括的に理解することが実現されつつある。また最近では、快と不快との感覚統合に関与する遺伝子の存在や、快と不快の連合学習の系についても報告されはじめている。我々は、*C. elegans* の温度走性の研究を通して、高等動物の感覚受容や学習・記憶機構の理解へ通じる分子生物学的知見を得たいと考えている。

脳の科学 25: 209-215, 2003

key words: *C. elegans*, nervous system, thermotaxis, sensory integration, learning and memory

I. はじめに

昨年度のノーベル医学・生理学賞が、Sydney Brenner 博士、H. Robert Horvitz 博士、John E. Sulston 博士に授与された。受賞理由は、線虫 *Caenorhabditis elegans* の器官発生と細胞死について、その遺伝学的制御の解明に多大なる貢献をしたというものである。今回初めて *C. elegans* の研究者がノーベル賞を受賞したことは、*C. ele-*

gans がモデル生物として世間に広く認知されたことの反映であるように思われ、同じ *C. elegans* を研究材料に扱っている者としてこの受賞は多大なる喜びであった。今となってはモデル生物として世界中で脚光を浴びている *C. elegans* ではあるが、ここまで *C. elegans* を用いた研究が発展したのは、Sydney Brenner 博士の存在をなくしては語ることができない。*C. elegans* が持つ特徴として、発生の全段階を通して体が透明なこと、自家受精で繁殖可能であること、世代時間が約3日と短いことなどの点が挙げられる。1960年代、Brenner 博士はこれらの特徴に着目し、発生や神経系などの高次生命現象を解明するためには、単純な構造を持ち遺伝学的手法が使用可能な *C. elegans* が最適だと考え、研究材料として *C. elegans* を使い始めた。それ以降の研究により、全細胞系譜、全神経回路網、全塩基配列が決定されるにしたがって、*C. elegans* は高次生命現象を個体、細胞、遺伝子レベルを一貫して研究でき

Love and hate in *C. elegans*: through the investigation of reception, integration, learning and memory of sensory signals.

*名古屋大学大学院理学研究科生命理学専攻分子神経生物学研究室

[〒464-8602 名古屋市中種区不老町]

Akiko Mohri and Ikue Mori: Laboratory of Molecular Neurobiology, Division of Biological Science, Graduate School of Science, Nagoya University, Fro-cho, Chikusa-ku, Nagoya, 464-8602 Japan.




図1 amphid 感覚ニューロンの神経終末

- a) *C. elegans* の頭部の模式図。A は頭側、P は尾側、D は背側、V は腹側を示す。
- b) 頭部先端部に位置する amphid 感覚子に存在する感覚ニューロンの終末部分を模式化したもの。amphid 感覚ニューロンは神経終末を外に向かって伸長させている。数種類の感覚ニューロンは繊毛を直接外部に露出している。温度受容ニューロン AFD の神経終末は外部に露出しておらず、sheath cell 内で微絨毛様構造を形成している。嗅覚ニューロン AWA, AWB, AWC の神経終末は直接外部に露出していないが、amphid の孔を通して間接的に外部に露出している。(文献2より改変)

る数少ないモデル生物として、確固たる地位を築いていった。

生命が生存するためには、刻々と変化する外部環境を察知し、その変化に対して素早い応答をすることが必須である。*C. elegans* は体長わずか1mm程度でありながら、外界の「快」「不快」を受容し、体内でその刺激のシグナルを統合、変換して、行動として出力することができる。本稿では、まず、温度走性を含めた *C. elegans* の感覚受容について概説し、次いで、最近になって報告がされ始めてきた、*C. elegans* にとっての「快」や「不快」を用いた感覚統合や学習・記憶の研究について、その内容の一部を紹介したい。

II. *C. elegans* の感覚：温度への応答に関与する神経回路と遺伝子

C. elegans は、成虫（雌雄同体）でわずか959個の体細胞でしか構成されていないにもかかわらず、基本的な組織である上皮、筋肉、消化管、神経系を全て兼ね備えている。神経系は302個のニューロンによって構成されており、*C. elegans* はその単純な神経系を駆使して、味、匂い、温度、高浸透圧、振動などの外部刺激に対して応答する。これらの刺激は、特定の感覚ニューロンで受容され、介在ニューロン、運動ニューロン、筋肉を介して、実際の行動に反映されと考えられている。外部刺激を受容する感覚ニューロンは、主に頭部に集中している。その中でも左右一対ず

表1 amphid 感覚ニューロン

感覚ニューロン		機能
AWA	化学受容 (嗅覚)	揮発性物質 (diacetyl, pyrazine, thiazole)
AWB	化学受容 (嗅覚)	揮発性物質に対する忌避
AWC	化学受容 (嗅覚)	揮発性物質 (benzaldehyde, butanone, isoamyl alcohol, thiazole)
AFD	温度受容	
ASE	化学受容 (味覚)	Na ⁺ , Cl ⁻ , cAMP, biotin, lysine, 産卵, 重金属イオン (Cd ²⁺ , Cu ²⁺) に対する忌避
ASG	化学受容 (味覚)	Na ⁺ , Cl ⁻ , cAMP, biotin, lysine
ASI	化学受容 (味覚)	Na ⁺ , Cl ⁻ , cAMP, biotin, lysine, 耐性幼虫フェロモン
ASJ	化学受容 (味覚)	耐性幼虫フェロモン
ASK	化学受容 (味覚)	lysine, 産卵
ADF	化学受容 (味覚)	耐性幼虫フェロモン
ASH		高浸透圧, 高濃度の揮発性物質, 重金属イオン (Cd ²⁺ , Cu ²⁺) に対する忌避
ADL		頭部先端部の接触による後退 octanol, 水溶性物質 (Cd ²⁺ , Cu ²⁺ も含む) に対する忌避




図2 野生型の温度走性

温度走性のアッセイは、中心から縁にかけて17℃～25℃の同心円状温度勾配を形成した9 cm シャーレ内の、餌のない寒天平板上でおこなう。写真は、餌を十分与えて飼育した野生型 *C. elegans* を約22℃である×印のところに置き、50分後の軌跡を撮影したものである。a) は17℃, b) は20℃, c) は25℃で飼育した野生型個体の軌跡をそれぞれ示す。いずれの飼育温度でも, *C. elegans* は過去の飼育温度付近に移動し, そこで等温線を描くように移動している。

つ存在する感覚子 amphid は最大規模であり, 12 種類, 計24個の感覚ニューロンにより構成されている¹⁹⁾。amphid に存在する各ニューロンの神経終末はそれぞれ特殊化した形態をしており, その機能と役割は, レーザーによる細胞殺傷実験および行動解析により明らかになっている^{2, 3, 10, 13, 14, 16)} (図1, 表1)。感覚ニューロンの軸索は, *C. elegans* の脳である nerve ring (神経環) に投射し

ており, そこで他のニューロンとシナプスを形成し, 下流の介在ニューロンへシグナルを伝達している。

我々の研究室では, 感覚受容の中でも特に温度受容機構に焦点を当てて研究している。*C. elegans* は, 餌の豊富な状態で飼育された場合, 餌のない温度勾配上で過去の飼育温度付近に移動し等温線を描く⁹⁾ (図2)。この性質は温度走性と呼




図3 温度走性を制御する神経回路モデル
矢印はシナプス結合、プラス(+)印は活性化シグナル、マイナス(-)印は不活性化シグナル、 \vdash 印はギャップ結合を示す。AFDは感覚ニューロン(温度受容細胞)、AIB, AIY, AIZ, RIA, RIBは介在ニューロン、RIMは運動ニューロン、Xはまだ未同定の温度受容細胞である。AIYは高い温度(U)への移動に、AIZは低い温度(D)への移動に関与する。AIBは低い温度(d)への移動にわずかに関与すると考えられる。(文献5より改変)

ばれ、*C. elegans*の行動の中でも神経可塑性が最も直接的に観察できる行動である。前述のように、*C. elegans*では、電子顕微鏡の切片解析により、全ニューロン間の化学シナプスとギャップ結合がすでに明らかにされている¹⁹⁾。当研究室の森らはこの利点を生かして、レーザー殺傷実験による温度走性に関与するニューロンおよび神経回路の同定を試みた結果、感覚ニューロンAFDが主要な温度受容ニューロンであること、介在ニューロンAIYの破壊によって個体は冷たい温度の方へ常に移動するようになること、介在ニューロンAIZの破壊によって個体は暖かい温度の方へ

常に移動するようになることなどを明らかにした。これらの結果から、AFD, AIY, AIZを含む、温度走性の成立に関与する主要な神経回路、およびその制御モデルが提唱された¹³⁾(図3)。その一方で、温度走性の分子機構を解明するために、多数の温度走性異常変異株が単離されており、その中の幾つかについてはすでに原因遺伝子がクローニングされている。その中の一部には、cGMP依存性チャネルの α サブユニットをコードする *tax-4* 遺伝子、OTX様転写因子をコードする *ttx-1* 遺伝子、calcineurinAサブユニットをコードする *tax-6* 遺伝子などがある^{11,12,17)}。そのいずれも、先に提唱された温度走性に関与すると予測された神経回路上のニューロンで発現・機能しており、温度走性に関与する神経回路制御モデルを支持するものとなっている。*C. elegans*の感覚行動の中で神経回路制御モデルが提唱されているのは、温度走性と機械刺激に対する反射に関してのみである^{5,13)}。したがって、温度走性に関与する遺伝子をさらにクローニングすることで、同じニューロン内での分子間の相互作用のみならず、ニューロンをまたいだ分子間の相互作用についても、新たな事実が解明されてくることが期待される。

Ⅲ. *C. elegans*における「快」と「不快」の研究から見てきたこと

*C. elegans*は、実際の飼育環境下では、外部環境から複数の「快」と「不快」を同時に受容していると予想される。*C. elegans*の感覚ニューロンは受容する刺激が分担されているため、*C. elegans*は、個別に受容された複数の刺激を下流の神経系で変換、統合し、最終的に筋肉を介して行動という形で出力していると思われる。それでは、*C. elegans*はどのような分子機構により複数の刺激を同時に処理しているのだろうか？この問いに対する答えを導き出すために、国立遺伝学研究所の石原らは、「快」物質である揮発性誘引物質の diacetyl と「不快」物質である水溶性忌避物質の銅イオンを用いて、快と不快の情報統合機構を調べるアッセイ系を確立した⁸⁾。このア

ッセイ系は、直径10cmの寒天平板の中央に寒天平板を2分割するように不快物質である銅イオンの線を引き、それを挟んで片方の区画に快物質のdiacetylを、もう片方の区画に*C. elegans*を置き、90分後にそれぞれの区画にいる個体数を数えるというものである。*C. elegans*が快物質のdiacetyl側に到達するためには、不快物質の銅イオンのバリアーを越えていかなければならないため、バリアーを越えてdiacetyl側にたどり着く個体の割合は、diacetylと銅イオンの濃度に依存する。石原らはこのアッセイ系を用いてスクリーニングをおこない、野生型よりも銅イオンのバリアーを越えにくい突然変異体 *hen-1* (*hesitation in crossing an aversive barrier*) を単離した。*hen-1* 変異体は、diacetyl または銅イオン単独に対しては、野生型と同様の誘引行動または忌避行動を示したが、diacetyl と銅イオンが同時に存在する場合には、野生型よりも銅イオンのバリアーを越えにくいという異常を示した。*hen-1* 変異の原因遺伝子をクローニングした結果、HEN-1タンパクは、LDLモチーフを持つ分泌タンパク質と高い相同性を持つことが明らかになった。また、*hen-1* 変異体は複数の学習・記憶のアッセイ系でも異常を示すことから、HEN-1分泌タンパク質は、感覚統合だけではなく、学習・記憶機構においても重要な役割を果たしていることが予想された。

また、東京大の佐伯らは、「快」物質である水溶性誘因物質のNaClと「不快」状態である飢餓を組み合わせたアッセイ系により、学習・記憶機構を解明しようと試みている¹⁵⁾。通常、餌(大腸菌)が豊富な状態で飼育している場合には、*C. elegans* はNaClに誘引されるが、NaClを含む餌のないプレートで4時間飼育した後は、NaClに誘引される個体が大幅に減少する。この現象は、NaClを含まない餌のないプレートで飼育した場合や、NaClを含んでも餌があるプレートで飼育した場合には観察されない。つまり、NaClと飢餓が同時に存在したときにのみ起こる、一種の連合学習ではないかと予想されている。さらに、NaClと飢餓との条件付けをした後は、NaClと同一の感覚ニューロンで受容される他の水溶性




図4 温度走性の予想概念図

誘引物質への化学走性も大幅に減少することから、飢餓によるNaClへの誘引の減少は、水溶性誘引物質への化学走性に関与するニューロンの誘引物質への応答が、細胞レベルで変化することによって引き起こされるのではないかと予想されている。

IV. 温度走性をモデルとした学習・記憶機構の解明への試み

1975年に発表されたHedgecockとRussellの論文では、温度走性という現象が報告されたが、同時に、温度への応答が飼育環境によって可逆的に変化することも報告された。例えば、*C. elegans*の飼育温度を16℃から25℃へ変化させた場合、25℃へ移した直後は、大部分の個体が16℃に近い低温領域に移動し25℃に近い高温領域に移動する個体は少ないが、25℃へ移してから約2～4時間後には、大部分の個体が25℃に近い高温領域に移動するようになった。この結果は、*C. elegans*がわずかな時間で新しい飼育温度を記憶したことを示唆する。また、餌がなく過密な悪環境で飼育した場合には、*C. elegans*は餌のない温度勾配上で過去の飼育温度から拡散するようになった⁹⁾。この行動変化は、飢餓状態または高濃度フェロモンあるいはその両方が原因であると予想され、*C. elegans*が飼育温度と悪環境とを関連付けて記憶していることを示唆する。

これらの結果から、温度走性の成立には、概念的に、①温度受容、②温度情報の記憶、③記憶さ

れている飼育時の温度と餌などの飼育状態との関連付け, ④温度勾配上で経時的に受容される現在の温度と記憶温度との相対評価, ⑤温度勾配上における温度評価の筋肉活動への反映, といった過程が正常に遂行されることが必須であると予想される(図4)。

先にも述べたが, 温度走性は *C. elegans* の行動の中でも神経可塑性が最も直接的に観察できる現象の一つである。我々は, 飼育温度や餌などの飼育環境を変化させた場合に起こる温度走性行動の変化は, *C. elegans* が新しい飼育環境を学習・記憶した結果ではないかと予想しており, 温度走性の成立機構を研究することで一般的な学習・記憶機構を解明していくことが可能であると考えている。すなわち, 飼育温度を条件刺激, 餌が豊富な状態を「快」, 餌がない状態を「不快」の無条件刺激とすると, 餌が豊富な状態で飼育された *C. elegans* が, 餌のない温度勾配上で過去の飼育温度に移動するのは, 任意の飼育温度と「快」(餌)を関連付けて記憶しているからであり, 餌のない状態で飼育された *C. elegans* が過去の飼育温度から拡散するのは, 任意の飼育温度と「不快」(飢餓)を関連付けて記憶しているからであると解釈することができる。

もし, 餌条件のみによって温度走性行動の変化を引き起こすことができるのであれば, 温度走性の成立に serotonin や octopamine などの神経伝達物質または神経ホルモンが関与している可能性が高い。*C. elegans* の咽頭の咀嚼行動や産卵行動などについての薬理学的解析により, serotonin は餌がある状態を, octopamine は餌がない状態をそれぞれ疑似することができることが明らかになっており, 両者は生体内で拮抗的に機能しているのではないかと予測されている^{1,7,10)}。この結果は, *C. elegans* では, 餌を食べたときには内在性の serotonin 濃度が上昇し, 餌がないときには内在性の octopamine 濃度が上昇する可能性を示唆する。現在のところ, serotonin と octopamine がどのように餌のシグナル伝達に関与しているのかについてはほとんど解明されていない。しかし, *Aplysia* や locust などでの薬理学的解析からも示唆されているように, serotonin や octo-

pamine が *C. elegans* においても神経可塑性に深く関与している可能性は高いと思われる。

V. おわりに

生命の高次機能を研究する場合, 行動(個体), 神経細胞(細胞), 遺伝子(分子)の各階層で多面的な解析が可能な *C. elegans* は, 非常に高いポテンシャルを持ったモデル生物である。特に, 神経系の発生・機能の研究においては, 全細胞系譜, 全神経回路網がすでに決定済みであることから, 哺乳類をはじめとする高等生物では非常に困難だと予想される任意の現象を, 神経回路レベルで包括的に解析することが可能である。*C. elegans* の神経系で最も特徴的なことは, ニューロンの数がわずかに302個であるにもかかわらず, 複数の「快」や「不快」を同時に受容・処理することが可能な上, 学習・記憶能力さえも持つことである。*C. elegans* の感覚ニューロンは数十種類しか存在しないため, 複数の異なる刺激を受容するためには, 同一のニューロンで複数のシグナルが交錯しながら, あるいは, 並列しながら伝達される必要がある。同時に, このようなシグナルの選択的伝達は神経回路レベルにも該当すると考えられる。様々な細胞の中でも, 神経細胞の化学的, および物理的な性質は, 進化の過程で最も保存されていると考えられるため, *C. elegans* の温度走性には, 高等動物の神経系を理解する上で重要なエッセンスが凝縮されているはずである。我々は, *C. elegans* の温度走性に関与する分子や神経細胞間のシグナル伝達系を解析することで, ヒトに至る高等動物の感覚受容や学習・記憶機構の理解へ通じる分子生物学的知見を得たいと考えている。

文 献

- 1) Avery, L., Horvitz, H.R.: Effects of starvation and neuroactive drugs on feeding in *Caenorhabditis elegans*. J. Exp. Zool., 253: 263-270, 1990.
- 2) Bargmann, C.I., Horvitz, H.R.: Chemosensory neurons with overlapping functions direct chemotaxis to multiple chemicals in *C. elegans*. Neuron, 7: 729-742, 1991.

- 3) Bargmann, C.I., Hartwig, E., Horvitz, H.R.: Odorant-selective genes and neurons mediate olfaction in *C. elegans*. *Cell*, 74: 515-527, 1993.
- 4) Bargmann, C.I.: Neurobiology of the *Caenorhabditis elegans* genome. *Science*, 282: 2028-2033, 1998.
- 5) Chalfie, M., Sulston, J.E., White, J.G. et al.: The neural circuit for touch sensitivity in *Caenorhabditis elegans*. *J. Neurosci.*, 5: 956-964, 1985.
- 6) Hedgecock, E.M., Russell, R.L.: Normal and mutant thermotaxis in the nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U.S.A.*, 72: 4061-4065, 1975.
- 7) Horvitz, H.R., Chalfie, M., Trent, C. et al.: Serotonin and octopamine in the nematode *Caenorhabditis elegans*. *Science*, 216: 1012-1014, 1982.
- 8) Ishihara, T., Iino, Y., Mohri, A. et al.: HEN-1, a secretory protein with an LDL receptor motif, regulates sensory integration and learning in *Caenorhabditis elegans*. *Cell*, 109: 639-649, 2002.
- 9) Jorgensen, E.M., Rankin, C.: Neural Plasticity. In: *C. elegans* II (ed. by Riddle, D.L., Blumenthal, Thomas, Meyer, B.J., Priess, J.R.), pp. 769-790, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1997.
- 10) Kaplan, J.M., Horvitz, H.R.: A dual mechanosensory and chemosensory neuron in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U.S.A.*, 90: 2227-2231, 1993.
- 11) Komatsu, H., Mori, I., Rhee, J.S. et al.: Mutations in a cyclic nucleotide-gated channel lead to abnormal thermosensation and chemosensation in *C. elegans*. *Neuron*, 17: 707-718, 1996.
- 12) Kuhara, A., Inada, H., Katsura, I. et al.: Negative regulation and gain control of sensory neurons by the *C. elegans* calcineurin TAX-6. *Neuron*, 33: 751-763, 2002.
- 13) Mori, I., Ohshima, Y.: Neural regulation of thermotaxis in *Caenorhabditis elegans*. *Nature*, 376: 344-348, 1995.
- 14) Perkins, L.A., Hedgecock, E.M., Thomson, J.N. et al.: Mutant sensory cilia in the nematode *Caenorhabditis elegans*. *Dev. Biol.*, 117: 456-487, 1986.
- 15) Saeki, S., Yamamoto, M., Iino, Y.: Plasticity of chemotaxis revealed by paired presentation of a chemoattractant and starvation in the nematode *Caenorhabditis elegans*. *J. Exp. Biol.*, 204: 1757-1764, 2001.
- 16) Sambongi, Y., Nagae, T., Liu, Y. et al.: Sensing of cadmium and copper ions by externally exposed ADL, ASE, and ASH neurons elicits avoidance response in *Caenorhabditis elegans*. *Neuroreport*, 10: 753-757, 1999.
- 17) Satterlee, J.S., Sasakura, H., Kuhara, A. et al.: Specification of thermosensory neuron fate in *C. elegans* requires *ttx-1*, a homolog of *otd/Otx*. *Neuron*, 31: 943-956, 2001.
- 18) Trent, C., Tsung, N., Horvitz, H.R.: Egg-laying defective mutants of the nematode *Caenorhabditis elegans*. *Genetics*, 104: 619-647, 1983.
- 19) White, J.G., Southgate, E., Thomson, J.N. et al.: The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philos. Trans. R. Soc. London Ser. B Biol. Sci.*, 314: 1-340, 1986.

レビュー

C. elegans の neuromodulator

C. elegans には、serotonin や octopamine 以外にも、acetylcholine, GABA (gamma-aminobutyric acid), dopamine, glutamate などの神経伝達物質が存在する。その中でも、dopamine, serotonin, octopamine は、野生型や変異体についての行動解析や薬理学的行動解析により、*C. elegans* でも neuromodulator として機能することが明らかになっている⁹⁾。他に neuromodulator として機能すると予想される物質に、FMRFamid (Phe-Met-Arg-Phe-amide) や神経ペプチドなどがあり、*C. elegans* のゲノム中にもこれらと相溶性が高い遺伝子が数十個存在することが予想されている⁴⁾。現段階では、これらのアミンやペプチドが神経伝達物質と機能しているのか、神経ホルモンとして機能しているのかははっきりしていない。しかし、もしこれらの neuromodulator の多くがホルモンとしても機能しているのであればなおさら、*C. elegans* が同時に複数のシグナルを処理するためにシナプスや神経筋接合部でシグナルの選択的伝達をおこなうことは必須条件であろう。